



---

Basic Research in Computer Science

BRICS RS-98-21 Power et al.: A Representation Result for Free Cocompletions

## A Representation Result for Free Cocompletions

John Power  
Gian Luca Cattani  
Glynn Winskel

BRICS Report Series

ISSN 0909-0878

RS-98-21

September 1998

**Copyright © 1998, BRICS, Department of Computer Science  
University of Aarhus. All rights reserved.**

**Reproduction of all or part of this work  
is permitted for educational or research use  
on condition that this copyright notice is  
included in any copy.**

**See back inner page for a list of recent BRICS Report Series publications.  
Copies may be obtained by contacting:**

**BRICS  
Department of Computer Science  
University of Aarhus  
Ny Munkegade, building 540  
DK-8000 Aarhus C  
Denmark  
Telephone: +45 8942 3360  
Telefax: +45 8942 3255  
Internet: BRICS@brics.dk**

**BRICS publications are in general accessible through the World Wide  
Web and anonymous FTP through these URLs:**

`http://www.brics.dk`  
`ftp://ftp.brics.dk`  
**This document in subdirectory RS/98/21/**

# A Representation Result for Free Cocompletions

John Power<sup>1,\*</sup>, Gian Luca Cattani<sup>2,†</sup>, Glynn Winskel<sup>3</sup>

<sup>1</sup>**LFCS**, University of Edinburgh, Scotland

<sup>2</sup>Computer Laboratory, University of Cambridge, England

<sup>3</sup> **BRICS**<sup>‡</sup>, University of Aarhus, Denmark

September 1998

## Abstract

Given a class  $F$  of weights, one can consider the construction that takes a small category  $\mathbb{C}$  to the free cocompletion of  $\mathbb{C}$  under weighted colimits, for which the weight lies in  $F$ . Provided these free  $F$ -cocompletions are small, this construction generates a 2-monad on **Cat**, or more generally on  $\mathcal{V}$ -**Cat** for monoidal biclosed complete and cocomplete  $\mathcal{V}$ . We develop the notion of a dense 2-monad on  $\mathcal{V}$ -**Cat** and characterise free  $F$ -cocompletions by dense  $KZ$ -monads on  $\mathcal{V}$ -**Cat**. We prove various corollaries about the structure of such 2-monads and their Kleisli 2-categories, as needed for the use of open maps in giving an axiomatic study of bisimulation in concurrency. This requires the introduction of the concept of a pseudo-commutativity for a strong 2-monad on a symmetric monoidal 2-category, and a characterisation of it in terms of structure on the Kleisli 2-category.

---

\*This work is supported by EPSRC grant GR/J84205: Frameworks for programming language semantics and logic.

<sup>†</sup>Research carried out while this author was at **BRICS**.

<sup>‡</sup>**B**asic **R**esearch in **C**omputer **S**cience, A Centre of the Danish National Research Foundation.



# 1 Introduction

Given a class of small categories  $S$ , one can consider the construction that takes a small category  $\mathbb{C}$  to the free cocompletion of  $\mathbb{C}$  under colimits of diagrams with shape in the class  $S$ . More subtly, and more naturally from the perspective of enriched category theory, for any monoidal biclosed complete and cocomplete  $\mathcal{V}$ , given a class  $F$  of weights, one can consider the construction that takes a small  $\mathcal{V}$ -category  $\mathbb{C}$  to the free cocompletion of  $\mathbb{C}$  under weighted colimits for which the weight lies in  $F$ . Provided that these  $S$ - or  $F$ -cocompletions are small, this construction generates a 2-monad on  $\mathbf{Cat}$ , or more generally on  $\mathcal{V}\text{-Cat}$ . We give an axiomatic characterisation of those 2-monads that arise in this way.

Although our characterisation and the results leading up to it hold for monoidal biclosed complete and cocomplete  $\mathcal{V}$ , we shall only state them for symmetric monoidal closed complete and cocomplete  $\mathcal{V}$ , because otherwise the swapping between  $\mathcal{V}$ -categories and  $V_t$ -categories, where  $V_t = \mathcal{V}$  but has the tensor product switched, becomes notationally tiresome. Some later results only hold for symmetric or even cartesian closed  $\mathcal{V}$ : where that is so, we state it explicitly.

The notion of  $KZ$ -monad was introduced as a property of 2-monads given by cocompletions under classes of colimits [11]. But it is not sufficient to characterise them. For instance, consider the 2-monad  $T$  on  $\mathbf{Cat}$  that sends every category to  $\mathbf{1}$ , the category with one object and one arrow only. It is a  $KZ$ -monad, but it is not given by a free cocompletion as there are many categories  $\mathbb{C}$  that do not embed fully into  $\mathbf{1}$ . So we need an additional condition. The condition we introduce is that of density of a 2-monad. We say that a 2-monad on  $\mathcal{V}\text{-Cat}$  is dense if both the unit  $\eta_{\mathbb{C}} : \mathbb{C} \rightarrow T\mathbb{C}$  and the  $\mathcal{V}$ -functor induced by the unit  $\tilde{\eta}_{\mathbb{C}} : T\mathbb{C} \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  are fully faithful, coherently with respect to the action of  $T$  on  $\mathcal{V}$ -functors. It follows that the coherence holds for  $\mathcal{V}$ -natural transformations too. The central result of this paper is the proof that a 2-monad on  $\mathcal{V}\text{-Cat}$  is a dense  $KZ$ -monad if and only if there is a class  $F$  of weights such that  $T$  is the 2-monad whose algebras are small  $F$ -cocomplete  $\mathcal{V}$ -categories.

En route to that result, we characterise dense  $KZ$ -monads as those 2-monads for which the  $\mathcal{V}$ -functors  $\eta_{\mathbb{C}}$  and  $\tilde{\eta}_{\mathbb{C}}$  are always fully faithful and such that, for any  $\mathcal{V}$ -functor  $H : \mathbb{C} \rightarrow T\mathbb{D}$ , the lifting of  $H$  to  $T\mathbb{C}$  given by the Kleisli construction is the restriction, up to coherent isomorphism, of  $\text{Lan}_{Y_{\mathbb{C}}}(\tilde{\eta}_{\mathbb{D}}H)$ , where  $Y_{\mathbb{C}} : \mathbb{C}^{\text{op}} \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  is the Yoneda embedding.

We further show consequences of the definition, or alternatively of the characterisation, of dense  $KZ$ -monads: it follows that  $T$  sends finite coproducts to finite products if finite coproducts exist coherently in each  $T\mathbb{C}$ ; in fact, the Kleisli 2-category has finite coproducts and they agree with finite products. Moreover, under a mild extra condition, every dense  $KZ$ -monad is pseudo-commutative, or equivalently, the Kleisli category is equivalent to a *Gray*-monoid, with structure induced by that in  $\mathcal{V}\text{-Cat}$ .

The structures developed here are not purely of abstract interest, but have been used in the study of concurrency. They are at the heart of attempts to model bisimulation by open maps [6, 3]. Given a monad  $T$  on  $\mathbf{Cat}$ , one may define a notion of open map within any category of the form  $T\mathbb{C}$ , with openness with respect to maps in  $\mathbb{C}$ . This notion of open map agrees with the notion of functional bisimulation, and can be extended, by considering spans of epimorphic open maps, to account for bisimulation in general as in [12]. One can prove, for a dense  $KZ$ -monad  $T$ , that all functors in the Kleisli category for  $T$  preserve open maps. The above structures allow one to account for the preservation of bisimulation by the various constructors, such as prefixing, nondeterministic sum, and a parallel operator, in a process algebra such as  $\mathbf{CCS}$ . See [3], which contains an early version of some of the results of this paper, for more detail.

The paper is organised as follows. In Section 2, we recall the definition of weighted colimit in a  $\mathcal{V}$ -category for monoidal biclosed complete and cocomplete  $\mathcal{V}$ . In Section 3, we motivate and define the notion of dense  $KZ$ -monad, and characterise it in terms of a condition on the Kleisli 2-category. In Section 4, we give our characterisation theorem. In Section 5, we develop the notion of a pseudo-commutativity for a strong 2-monad on a symmetric monoidal 2-category, and we characterise it in terms of a *Gray*-monoid structure on the Kleisli 2-category. Finally, in Section 6, we deduce further properties of dense  $KZ$ -monads, requiring some further conditions.

## 2 Weighted colimits and free cocompletions

In this section, we recall some definitions and results associated with weighted colimits and free cocompletions. The standard reference for these, in the case of enrichment over symmetric monoidal closed  $\mathcal{V}$ , is Kelly's book [8]. For the case of monoidal biclosed  $\mathcal{V}$ , some more delicacy is needed, and the definitions for such  $\mathcal{V}$ , more generally for enrichment in a biclosed bicategory  $\mathcal{W}$ , are

in [4]. For ease of exposition, we shall write in terms of symmetric  $\mathcal{V}$ .

**2.1 Definition** A *weight* is a small  $\mathcal{V}$ -category  $\mathbb{D}$  together with a  $\mathcal{V}$ -functor

$$f : \mathbb{D}^{\text{op}} \longrightarrow \mathcal{V} .$$

**2.2 Definition** Given a weight  $f : \mathbb{D}^{\text{op}} \longrightarrow \mathcal{V}$  and a  $\mathcal{V}$ -functor  $g : \mathbb{D} \longrightarrow \mathbb{C}$ , an  *$f$ -weighted colimit of  $g$*  is an object  $\text{colim}(f, g)$  of  $\mathbb{C}$  together with, for each object  $X$  of  $\mathbb{C}$ , an isomorphism between  $[\mathbb{D}^{\text{op}}, \mathcal{V}](f-, \mathbb{C}(g-, X))$  and  $\mathbb{C}(\text{colim}(f, g), X)$ ,  $\mathcal{V}$ -naturally in  $X$ .

This definition can equally be expressed as asking for the existence of a colimiting cylinder [8] (rather than cone), such that composition with that cylinder yields the above natural isomorphism for every  $X$ . Here, a cylinder is just an object of  $[\mathbb{D}^{\text{op}}, \mathcal{V}](f-, \mathbb{C}(g-, X))$ , extending the notion of cone, which is essentially the case where  $f$  is the constant at 1: it is not quite that simple, mainly because for arbitrary  $\mathcal{V}$ , the unit need not be the terminal object, and consequently constant  $\mathcal{V}$ -functors need not exist. See [8] for more detail on that. Nevertheless, weighted colimits do include all conical colimits. They also include all tensors, which you obtain by considering  $\mathbb{D} = 1$ . For the case of  $\mathcal{V} = \mathbf{Set}$ , i.e., for ordinary categories, tensors are copowers, so the concept of weighted colimit loses much of its force. But that is not true for  $\mathcal{V}$ -categories in general, for instance for  $\mathcal{V} = \mathbf{Poset}$  or  $\mathcal{V} = \mathbf{Cat}$ . And even in the case of ordinary categories, our characterisation here of free cocompletions under weighted colimits cannot be expressed simply in terms of free cocompletions under conical colimits, as we shall see.

**2.3 Definition** Given a class  $F$  of weights, a locally small  $\mathcal{V}$ -category with all  $f$ -weighted colimits for all  $f$  in  $F$  is called  *$F$ -cocomplete*. A  $\mathcal{V}$ -functor between  $F$ -cocomplete  $\mathcal{V}$ -categories that preserves all  $f$ -weighted colimits for all  $f$  in  $F$  is called  *$F$ -cocontinuous*.

**2.4 Definition** Given a class  $F$  of weights, a *2-monad for  $F$ -cocompletions* is a 2-monad  $T$  on  $\mathcal{V}\text{-Cat}$  for which, for every small  $\mathcal{V}$ -category  $\mathbb{C}$ , the unit  $\eta_{\mathbb{C}} : \mathbb{C} \longrightarrow T\mathbb{C}$  exhibits  $T\mathbb{C}$  as the free  $F$ -cocompletion of  $\mathbb{C}$ , i.e., for any locally small  $F$ -cocomplete  $\mathcal{V}$ -category  $\mathbb{D}$ , composition with  $\eta_{\mathbb{C}}$  gives an equivalence between the category of  $F$ -cocontinuous  $\mathcal{V}$ -functors from  $T\mathbb{C}$  to  $\mathbb{D}$  and that of  $\mathcal{V}$ -functors from  $\mathbb{C}$  to  $\mathbb{D}$ .

Letting  $F\text{-Coc}$  denote the 2-category of small  $F$ -cocomplete  $\mathcal{V}$ -categories,  $F$ -cocontinuous  $\mathcal{V}$ -functors, and all  $\mathcal{V}$ -natural transformations, an immediate consequence of the definition is

**2.5 Corollary** If  $T$  is a 2-monad for  $F$ -cocompletions, then  $T$  provides a left biadjoint to the forgetful 2-functor  $U : F\text{-Coc} \rightarrow \mathcal{V}\text{-Cat}$ .

The corollary does not have a converse. For instance, if  $\mathcal{V} = \mathbf{Set}$  and  $F$  consists of all possible weights, then  $F\text{-Coc}$  is biequivalent to the 2-category of complete lattices, as every small cocomplete category is a preordered set; and a left biadjoint to  $U$  is not what we mean by the free cocompletion of a small category. If  $F$  is the class of all weights, then there is no 2-monad for free cocompletions, precisely because the free cocompletion of a small category is not small. However, it follows from the corollary that if it exists, a 2-monad for  $F$ -cocompletions is unique up to equivalence.

Note that we have been careful to consider  $F$ -cocontinuous  $\mathcal{V}$ -functors, not  $\mathcal{V}$ -functors that strictly preserve  $F$ -weighted colimits. The situation we consider is the natural one, but we remark that our results here do not restrict well, in that the 2-monads we consider do not act as free  $F$ -cocompletions with respect to  $\mathcal{V}$ -functors that strictly preserve  $F$ -weighted colimits. For an account of some of the relevant issues, and for our 2-categorical notation, see [2].

The main result of this paper is a characterisation of 2-monads for  $F$ -cocompletions. The central background result we use, shown for symmetric  $\mathcal{V}$  in [8], extending routinely to non-symmetric  $\mathcal{V}$  by the work of [4], is

**2.6 Theorem** The free  $F$ -cocompletion of a small  $\mathcal{V}$ -category  $\mathbb{C}$  is given by the closure of the representables in  $[\mathbb{C}^{\text{op}}, \mathcal{V}]$  under  $F$ -weighted colimits.

### 3 Dense $KZ$ -monads

In this section, we motivate and define the notion of dense  $KZ$ -monad, and we characterise them in terms of their Kleisli categories. In order to fix notation, we recall

**3.1 Definition** A 2-monad on a 2-category  $\mathcal{A}$  consists of a 2-functor

$$T : \mathcal{A} \rightarrow \mathcal{A},$$



together with 2-natural transformations  $\eta : id \Rightarrow T$  and  $\mu : T^2 \Rightarrow T$ , satisfying the usual three axioms treated as axioms about 2-natural transformations rather than ordinary natural transformations.

**3.2 Definition** Given a 2-monad  $T$  on  $\mathcal{A}$ , the *Kleisli 2-category* has as objects all objects  $C$  of  $\mathcal{A}$ , and arrows those arrows  $H : TC \rightarrow TD$  that respect  $\mu_C$  and  $\mu_D$ , with composition given by composition in  $\mathcal{A}$ . An arrow in  $Kl(T)$  may equivalently be described as any arrow in  $\mathcal{A}$  from  $C$  to  $TD$ . The 2-cells are those 2-cells in  $\mathcal{A}$  that respect  $\mu$ .

Considerable detail of 2-monads and the category theoretic constructions associated with them appears in [9].

**3.3 Definition** A *KZ-monad* is a 2-monad for which  $\mu : T^2 \Rightarrow T$  is left adjoint to  $\eta T$ . It is equivalent to ask that  $\mu$  be right adjoint to  $T\eta$  (see [11]).

The notion of *KZ-monad* was introduced to study particular features of 2-monads given by free cocompletions under classes of colimits [11]. But they do not characterise such free cocompletions, as the following example shows.

**3.4 Example** Consider the 2-monad on **Cat** that sends every category to the one object one arrow category **1**. It is a *KZ-monad* trivially, but it does not give free cocompletions under a class of colimits because **C** typically is not a full subcategory of **1**.

So we need to consider a condition that is stronger than that of being a *KZ-monad*.

### 3.5 Notation

- Given a 2-monad  $T$  on  $\mathcal{V}\text{-Cat}$ , let  $\tilde{\eta}_C : TC \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  denote the  $\mathcal{V}$ -functor that sends an object  $X$  to the  $\mathcal{V}$ -functor

$$TC(\eta_C -, X) : \mathbb{C}^{\text{op}} \rightarrow \mathcal{V} .$$

- Given  $\mathcal{V}$ -functors  $H : \mathcal{C} \rightarrow \mathcal{D}$  and  $J : \mathcal{C} \rightarrow \mathcal{C}'$ , the *left Kan extension of  $H$  along  $J$*  is given by a  $\mathcal{V}$ -functor  $\text{Lan}_J H : \mathcal{C}' \rightarrow \mathcal{D}$  and a  $\mathcal{V}$ -natural transformation  $\alpha : H \Rightarrow (\text{Lan}_J H)J$  that is universal among such  $\mathcal{V}$ -natural transformations, i.e., given any  $\mathcal{V}$ -functor  $K : \mathcal{C}' \rightarrow \mathcal{D}$  and any  $\mathcal{V}$ -natural transformation  $\beta : H \Rightarrow KJ$ , there exists a unique  $\mathcal{V}$ -natural transformation  $\gamma : \text{Lan}_J H \Rightarrow K$  making the evident triangle commute.

If it exists, a left Kan extension is unique up to coherent isomorphism. If  $J$  is fully faithful and a left Kan extension exists, then  $\alpha$  is necessarily an isomorphism. The left Kan extension always exists if  $\mathcal{C}$  is a small  $\mathcal{V}$ -category and  $\mathcal{D}$  is cocomplete (see [8] for more detail on left Kan extensions).

**3.6 Definition** A 2-monad  $T$  on  $\mathcal{V}\text{-Cat}$  is *dense* if for every small  $\mathcal{V}$ -category  $\mathbb{C}$ , the  $\mathcal{V}$ -functors  $\eta_{\mathbb{C}} : \mathbb{C} \rightarrow T\mathbb{C}$  and  $\tilde{\eta}_{\mathbb{C}} : T\mathbb{C} \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  are fully faithful, and for any  $H : \mathbb{C} \rightarrow \mathbb{D}$ , the  $\mathcal{V}$ -functor

$$\text{Lan}_{Y_{\mathbb{C}}}(Y_{\mathbb{D}}H) : [\mathbb{C}^{\text{op}}, \mathcal{V}] \rightarrow [\mathbb{D}^{\text{op}}, \mathcal{V}]$$

restricts to  $TH : T\mathbb{C} \rightarrow T\mathbb{D}$  up to coherent isomorphism, where  $Y_{\mathbb{C}} : \mathbb{C} \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  is the Yoneda embedding.

It follows from our conditions that the behaviour of  $T$  on 2-cells is given by the restriction of that determined by left Kan extension on  $[\mathbb{C}^{\text{op}}, \mathcal{V}]$ , cf the proof of Corollary 3.8.

For 2-monads given by free cocompletions,  $\eta_{\mathbb{C}} : \mathbb{C} \rightarrow T\mathbb{C}$  and  $\tilde{\eta}_{\mathbb{C}} : T\mathbb{C} \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  are fully faithful by Theorem 2.6, and for every  $\mathcal{V}$ -functor  $H : \mathbb{C} \rightarrow \mathbb{D}$ , the  $\mathcal{V}$ -functor  $\text{Lan}_{Y_{\mathbb{C}}}(Y_{\mathbb{D}}H) : [\mathbb{C}^{\text{op}}, \mathcal{V}] \rightarrow [\mathbb{D}^{\text{op}}, \mathcal{V}]$  restricts to  $TH : T\mathbb{C} \rightarrow T\mathbb{D}$  up to coherent isomorphism (see [8]).

**3.7 Theorem** Let  $T$  be a 2-monad on  $\mathcal{V}\text{-Cat}$  for which  $\eta_{\mathbb{C}} : \mathbb{C} \rightarrow T\mathbb{C}$  and  $\tilde{\eta}_{\mathbb{C}} : T\mathbb{C} \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  are fully faithful for every  $\mathbb{C}$ . Then  $T$  is a dense  $KZ$ -monad if and only if every  $\mathcal{V}$ -functor  $H : T\mathbb{C} \rightarrow T\mathbb{D}$  in  $Kl(T)$  is the restriction of  $\text{Lan}_{Y_{\mathbb{C}}}(\tilde{\eta}_{\mathbb{D}}H\eta_{\mathbb{C}}) : [\mathbb{C}^{\text{op}}, \mathcal{V}] \rightarrow [\mathbb{D}^{\text{op}}, \mathcal{V}]$  up to coherent isomorphism.

**Proof** Suppose  $T$  is dense and  $KZ$ , and let  $H : T\mathbb{C} \rightarrow T\mathbb{D}$  be a  $\mathcal{V}$ -functor in  $Kl(T)$ . Then  $H = \mu_{\mathbb{D}}K$  where  $K = H\eta_{\mathbb{C}} : \mathbb{C} \rightarrow T\mathbb{D}$ . Using the density condition applied to  $K$  and the definition of  $KZ$ -monad, and the fact that left Kan extensions into cocomplete  $\mathcal{V}$ -categories (such as  $[\mathbb{D}^{\text{op}}, \mathcal{V}]$ ) are colimits, so are preserved by  $\mathcal{V}$ -functors with right  $\mathcal{V}$ -adjoints, gives the result.

For the converse, given  $L : \mathbb{C} \rightarrow \mathbb{D}$ , let  $H = TL$ . By  $\mathcal{V}$ -naturality of  $\eta$  and since  $\tilde{\eta}_{\mathbb{D}}\eta_{\mathbb{D}}$  is isomorphic to  $Y_{\mathbb{D}} : \mathbb{D} \rightarrow [\mathbb{D}^{\text{op}}, \mathcal{V}]$  by fully faithfulness of  $\eta_{\mathbb{D}}$ , it follows that  $\tilde{\eta}_{\mathbb{D}}H\eta_{\mathbb{C}} = Y_{\mathbb{D}}L$ , so  $T$  is dense.

To see that  $T$  is  $KZ$ , first observe that  $\mu_{\mathbb{C}} : T^2\mathbb{C} \rightarrow T\mathbb{C}$  is a  $\mathcal{V}$ -functor in  $Kl(T)$ . So, up to isomorphism,  $\mu_{\mathbb{C}}$  is the restriction of

$$\text{Lan}_{Y_{T\mathbb{C}}}(\tilde{\eta}_{\mathbb{C}}\mu_{\mathbb{C}}\eta_{T\mathbb{C}}) = \text{Lan}_{Y_{T\mathbb{C}}}\tilde{\eta}_{\mathbb{C}} .$$

But, by fully faithfulness of  $\tilde{\eta}_{\mathbb{C}}$ , the  $\mathcal{V}$ -functor  $\eta_{T\mathbb{C}} : T\mathbb{C} \rightarrow T^2\mathbb{C}$  is the restriction of the  $\mathcal{V}$ -functor sending  $K\epsilon[\mathbb{C}^{\text{op}}, \mathcal{V}]$  to  $[\mathbb{C}^{\text{op}}, \mathcal{V}](\tilde{\eta}_{\mathbb{C}}-, K)$ , but this latter  $\mathcal{V}$ -functor is the right  $\mathcal{V}$ -adjoint of  $\text{Lan}_{Y_{T\mathbb{C}}}\tilde{\eta}_{\mathbb{C}}$ . Since  $\tilde{\eta}_{\mathbb{C}}$  and  $\tilde{\eta}_{T\mathbb{C}}$  are both fully faithful, it follows that  $\mu_{\mathbb{C}}$  is left  $\mathcal{V}$ -adjoint to  $\eta_{T\mathbb{C}}$ . ■

**3.8 Corollary** For any dense  $KZ$ -monad on  $\mathcal{V}\text{-Cat}$ , every  $H$  in  $Kl(T)$  is a left Kan extension of  $H\eta_{\mathbb{C}} : \mathbb{C} \rightarrow T\mathbb{D}$  along  $\eta_{\mathbb{C}} : \mathbb{C} \rightarrow T\mathbb{C}$ .

**Proof** Given any  $K : T\mathbb{C} \rightarrow T\mathbb{D}$ , it follows by fully faithfulness of  $\tilde{\eta}_{\mathbb{C}} : T\mathbb{C} \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  that  $\tilde{\eta}_{\mathbb{D}}K$  is isomorphic to  $\text{Lan}_{\tilde{\eta}_{\mathbb{C}}}(\tilde{\eta}_{\mathbb{D}}K)\tilde{\eta}_{\mathbb{C}}$ . Moreover, since  $\eta_{\mathbb{C}}$  is fully faithful, the Yoneda embedding  $Y_{\mathbb{C}} : \mathbb{C} \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$  is isomorphic to  $\tilde{\eta}_{\mathbb{C}}\eta_{\mathbb{C}}$ . So any  $\mathcal{V}$ -natural transformation  $\alpha : H\eta_{\mathbb{C}} \Rightarrow K\eta_{\mathbb{C}}$ , induces a  $\mathcal{V}$ -natural transformation from  $\tilde{\eta}_{\mathbb{D}}H\eta_{\mathbb{C}} : \mathbb{C} \rightarrow [\mathbb{D}^{\text{op}}, \mathcal{V}]$  to  $\text{Lan}_{\tilde{\eta}_{\mathbb{C}}}(\tilde{\eta}_{\mathbb{D}}K)Y_{\mathbb{C}}$ , hence by definition of left Kan extension, a  $\mathcal{V}$ -natural transformation

$$\bar{\alpha} : \text{Lan}_{Y_{\mathbb{C}}}(\tilde{\eta}_{\mathbb{D}}H\eta_{\mathbb{C}}) \Rightarrow \text{Lan}_{\tilde{\eta}_{\mathbb{C}}}(\tilde{\eta}_{\mathbb{D}}K) .$$

The result follows immediately from Theorem 3.7, the commutativity up to coherent isomorphism for  $K$ , and fully faithfulness of  $\tilde{\eta}_{\mathbb{D}}$ . ■

**3.9 Corollary** Given a dense  $KZ$ -monad  $T$  on  $\mathcal{V}\text{-Cat}$ , for every  $\mathbb{C}$ ,  $\mu_{\mathbb{C}}$  is the left Kan extension of  $id : \mathbb{C} \rightarrow \mathbb{C}$  along  $\eta_{T\mathbb{C}} : T\mathbb{C} \rightarrow T^2\mathbb{C}$ . Moreover, it is the restriction of  $[\eta-, \mathcal{V}] : [T\mathbb{C}^{\text{op}}, \mathcal{V}] \rightarrow [\mathbb{C}^{\text{op}}, \mathcal{V}]$ .

## 4 The representation theorem

For a dense  $KZ$ -monad  $T$ , it is not true that there exists a class  $S$  of small  $\mathcal{V}$ -categories such that  $T$  is the 2-monad for  $S$ -cocompletions. This is the case even for  $\mathcal{V} = \mathbf{Set}$ , i.e., even for ordinary categories.

**4.1 Example** Let  $T$  be the 2-monad on  $\mathbf{Cat}$  such that  $T\mathbf{0} = \mathbf{0}$ , where  $\mathbf{0}$  is the empty category and  $T\mathbb{C}$  is given by freely adding an initial object to  $\mathbb{C}$ . It is routine to check the 2-monad axioms and to verify that  $T$  is dense and  $KZ$ . However, it is not a free cocompletion. Suppose otherwise. Let  $S$  be a class of small categories such that  $T\mathbb{C}$  is the free  $S$ -cocompletion of  $\mathbb{C}$ . Since  $T\mathbf{0}$  is empty, every category in  $S$  must be nonempty. Now consider the transfinite construction of the free  $S$ -cocompletion. Since each category

in  $S$  is nonempty, every colimit that is introduced has a map into it from an already existing object, so ultimately from an object in  $\mathbb{C}$ . So at no point in the transfinite induction do we introduce an initial object, a contradiction.

We now state our representation theorem. The above counterexample means we need to consider weighted colimits rather than restricting ourselves to conical ones. Moreover, we need a little caution about the precise statement here, because the class  $F$  is typically not small. So the assumption that  $T$  is a 2-monad is important to ensure that the free  $F$ -cocompletion of a small  $\mathcal{V}$ -category is small.

**4.2 Theorem** A 2-monad  $T$  on  $\mathcal{V}\text{-Cat}$  is a dense  $KZ$ -monad if and only if there is a class  $F$  of weights for which  $T$  is the 2-monad for  $F$ -cocompletions.

**Proof** The forward direction of this largely follows immediately from Theorem 2.6. It is routine to verify from the universal property and construction in [8] that the 2-monad is a  $KZ$ -monad.

For the converse, given  $T$ , define  $F$  to be the disjoint union over all small  $\mathcal{V}$ -categories  $D$  of  $\{f : \mathbb{D}^{\text{op}} \rightarrow \mathcal{V} \mid f \in T\mathbb{D}\}$ . We must show for every small  $\mathcal{V}$ -category  $\mathbb{C}$ , that  $T\mathbb{C}$  is the closure in  $[\mathbb{C}^{\text{op}}, \mathcal{V}]$  of the representables under colimits of diagrams with weight in  $F$ . Trivially,  $T\mathbb{C}$  is contained in the latter  $\mathcal{V}$ -category, as every object of  $T\mathbb{C}$  lies in  $F$ , and is the colimit of representables determined by itself as a weight. So it remains to show for any small  $\mathcal{V}$ -category  $\mathbb{D}$ , any  $X$  in  $T\mathbb{D}$ , and any  $g : \mathbb{D} \rightarrow T\mathbb{C}$ , that the weighted colimit  $\text{colim}(\tilde{\eta}_{\mathbb{D}}X, \tilde{\eta}_{\mathbb{C}}g)$  in  $[\mathbb{C}^{\text{op}}, \mathcal{V}]$  lies in  $T\mathbb{C}$ . But this colimit is  $(\text{Lan}_{Y_{\mathbb{D}}}(\tilde{\eta}_{\mathbb{C}}g))\tilde{\eta}_{\mathbb{D}}X$ , and  $\text{Lan}_{Y_{\mathbb{D}}}(\tilde{\eta}_{\mathbb{C}}g)$  restricts to the lifting of  $g$ . So the colimit does lie in  $T\mathbb{C}$  and is given by the lifting of  $g$ . ■

We can gain a little more out of this with care. Given a small class  $F$  of weights, such as the class of weights for all finite colimits, one can consider the 2-monad  $T_F$  for  $F$ -cocompletions. One can then use the construction of the Theorem to obtain another class of weights  $F'$  such that  $T$  is the 2-monad for  $F'$ -cocompletions. But by construction,  $F'$  will be large. But we can characterise  $F'$  directly in terms of  $F$ . Albert and Kelly [1] defined a notion of closed class of weights.

**4.3 Definition** A class  $F$  of weights is *closed* if for any weight  $f : \mathbb{D}^{\text{op}} \rightarrow \mathcal{V}$ , if every  $F$ -cocomplete  $\mathcal{V}$ -category has  $f$ -weighted colimits, and if every  $F$ -cocontinuous  $\mathcal{V}$ -functor preserves  $f$ -weighted colimits, then  $f$  lies in  $F$ .

Any class of weights  $F$  has a closure  $\bar{F}$ . For instance, if  $F$  includes weights for finite products and equalisers, then  $\bar{F}$  includes weights for all finite limits. Albert and Kelly showed

**4.4 Proposition** A weight  $f : \mathbb{D}^{\text{op}} \rightarrow \mathcal{V}$  is in  $\bar{F}$  if and only if it is in the closure of the representables in  $[\mathbb{D}^{\text{op}}, \mathcal{V}]$  under  $F$ -weighted colimits.

So, it follows from the construction of the Theorem that we have

**4.5 Corollary** The class of weights given by the construction of the Theorem is always closed. And for any class  $F$  of weights for which  $F$ -cocompletions of small  $\mathcal{V}$ -categories are small, the class of weights given by the construction of the Theorem is the closure  $\bar{F}$  of  $F$ .

## 5 Pseudo-commutativity

In order to study the properties of dense  $KZ$ -monads, we need a notion of pseudo-commutativity of a strong 2-monad on a symmetric monoidal 2-category, and we need a result characterising pseudo-commutative 2-monads in terms of the Kleisli construction. So we develop such definitions and results in this section. A notion of pseudo-commutativity appeared in Kelly's [7], but was not developed in general. We have the same notion but more compact axioms.

**5.1 Definition** Let  $\mathcal{V}$  be a symmetric monoidal 2-category. A *pseudo-commutativity* for a strong 2-monad  $T$  on  $\mathcal{V}$  consists of, for every pair of objects  $(C, D)$ , an isomorphic 2-cell  $\chi_{C,D}$  between the two maps from  $TC \otimes TD$  to  $T(C \otimes D)$  induced by the strength, such that  $\chi$  is natural in  $C$  and  $D$  and satisfies three coherence conditions:  $T(c_{C,D})\chi_{C,D} = (\chi_{D,C})^{-1}c_{TC,TD}$ , coherence with respect to  $\mu_C$ , and  $\chi_{C,D}(\eta_C \otimes id_{TD}) = id$  (coherence for  $D$  being a consequence). A *pseudo-commutative monad* is a strong 2-monad together with a pseudo-commutativity.

In general, a strong monad  $T$  on a symmetric monoidal category is commutative if and only if  $Kl(T)$  is symmetric monoidal, with tensor product given by  $T(C \otimes D)$ . We generalise that result. Modulo a straightforward generalisation of the usual coherence relating monoidal and strict monoidal categories, the notion we need here is that of *Gray-monoid*.

The category  $\mathbf{2-Cat}$  has a symmetric monoidal closed structure for which the closed structure is given by the 2-category  $Ps(\mathbb{C}, \mathbb{D})$  of 2-functors from  $\mathbb{C}$  to  $\mathbb{D}$  and pseudo-natural transformations between them. The category  $\mathbf{2-Cat}$  together with this symmetric monoidal closed structure is called *Gray* [5].

**5.2 Definition** A *Gray-monoid* is a monoid in the symmetric monoidal closed category *Gray*, i.e., a 2-category  $\mathcal{A}$  together with a 2-functor

$$\cdot : \mathcal{A} \otimes \mathcal{A} \longrightarrow \mathcal{A} ,$$

subject to the monoid laws.

Spelling this out, we have

**5.3 Proposition** A *Gray-monoid* consists of

- a 2-category  $\mathcal{A}$ ,
- for each object  $X$  of  $\mathcal{A}$ , 2-functors  $h_X : \mathcal{A} \longrightarrow \mathcal{A}$  and  $k_X : \mathcal{A} \longrightarrow \mathcal{A}$ , such that for every pair of objects  $(X, Y)$ ,  $h_X Y = k_Y X$ , which we denote by  $X \otimes Y$ ,
- for each pair of maps  $f : X \longrightarrow X'$  and  $g : Y \longrightarrow Y'$ , a natural isomorphism between the two induced maps from  $X \otimes Y$  to  $X' \otimes Y'$

such that the isomorphisms respect the 2-functorial structure of  $h_X$  and  $k_X$ , and all the data respects associativity and units for  $\otimes$ .

**5.4 Definition** A *symmetry* for a *Gray-monoid* consists of, for each  $X$  and  $Y$ , an isomorphism  $c : X \otimes Y \longrightarrow Y \otimes X$ , such that  $c^2 = id$  and  $(id_Y \otimes c_{X,Z})(c_{X,Y} \otimes id_Z) = c_{X,Y \otimes Z}$ , and such that  $\mathbb{C}$  respects all the *Gray*-structure. A *symmetric Gray-monoid* is a *Gray-monoid* together with a symmetry on it.

We want to characterise a pseudo-commutativity of a strong monad  $T$  by the structure of a *Gray-monoid* on  $Kl(T)$ . That involves some delicate coherence. So we first recall a characterisation of a strength for a monad on a symmetric monoidal category in terms of structure on the Kleisli category [13]. A strict premonoidal category is exactly a *Gray-monoid* but with no two-dimensional data. In the following result, the notion of strict symmetric premonoidal functor is not entirely obvious, as it includes the axiom

that the functor preserves central maps, as we shall define in our setting shortly.

**5.5 Theorem** Let  $\mathcal{V}$  be a symmetric monoidal category, and let  $T$  be a monad on it. Then, to give a strength for  $T$  is to give a symmetric premonoidal structure on  $Kl(T)$  such that the canonical functor  $J : \mathcal{V} \rightarrow Kl(T)$  is a strict symmetric premonoidal functor.

We generalise the constructions of this result.

**5.6 Definition** A 1-cell  $f : X \rightarrow X'$  in a *Gray*-monoid is *central* if for every 1-cell  $g : Y \rightarrow Y'$ , the isomorphic 2-cell between  $(X' \otimes g)(f \otimes Y)$  and  $(f \otimes Y')(X \otimes g)$  is the identity, and the dual.

**5.7 Theorem** Given a strong 2-monad  $T$  on a strict symmetric monoidal 2-category  $\mathcal{V}$ , to give a pseudo-commutativity for  $T$  is to give  $Kl(T)$  the structure of a symmetric *Gray*-monoid such that the canonical 2-functor  $J : \mathcal{V} \rightarrow Kl(T)$  strictly preserves the strict symmetric monoidal structure, with each 1-cell in  $\mathcal{V}$  sent to a central 1-cell in  $Kl(T)$ .

**Proof** Given a pseudo-commutativity for  $T$ , and given  $f : TX \rightarrow TX'$  and  $g : TY \rightarrow TY'$  in  $Kl(T)$ , a study of the construction of the two composites from  $T(X \otimes Y)$  to  $T(X' \otimes Y')$  in  $Kl(T)$  yields the desired isomorphism, and the coherence laws follow directly from the coherence for pseudo-commutativity.

Conversely, to obtain a pseudo-commutativity from a symmetric *Gray*-monoid structure on  $Kl(T)$ , consider the components  $\epsilon_X : T^2X \rightarrow TX$  and  $\epsilon_Y : T^2Y \rightarrow TY$  of the counit  $\epsilon$  of the adjunction between  $\mathcal{V}$  and  $Kl(T)$ , and use the isomorphism determined by them. The coherence follows directly.

It is routine to verify that these constructions are mutually inverse.

■

We have glossed over a coherence issue here by assuming that  $\mathcal{V}$  is strict symmetric monoidal. If  $\mathcal{V}$  is symmetric monoidal, we cannot obtain a *Gray*-monoid structure on  $Kl(T)$  because the monoidal structure is not strict on the objects of  $Kl(T)$ . The most natural structure to consider in this regard is that of a monoidal bicategory, or a one object tricategory [5]. There is a coherence theorem that says every monoidal bicategory is equivalent to a *Gray*-monoid. But although the structure on  $Kl(T)$  is more general than that of a *Gray*-monoid, it is not as complicated as the structure of a

monoidal bicategory, as the monoidal structure of  $Kl(T)$  involves coherent isomorphisms rather than coherent equivalences, and several commutativities hold here but do not hold in an arbitrary monoidal bicategory. The structure does not seem important enough to merit its own name, but we can at least say what structure  $Kl(T)$  has.

**5.8 Theorem** To give a pseudo-commutativity for a strong 2-monad  $T$  on a symmetric monoidal 2-category  $\mathcal{V}$  is equivalent to the assertion that the Kleisli 2-category  $Kl(T)$  has the following structure:

- for each object  $X$ , 2-functors  $h_X : Kl(T) \rightarrow Kl(T)$  and  $k_X : Kl(T) \rightarrow Kl(T)$  such that  $h_X Y = k_Y X$ , extending  $\otimes$  on  $\mathcal{V}$ ,
- for each pair of maps  $f : TX \rightarrow TX'$  and  $g : TY \rightarrow TY'$  in  $Kl(T)$ , a natural isomorphism between the two induced maps from  $T(X \otimes Y)$  to  $T(X' \otimes Y')$ , extending the commutativity on  $\mathcal{V}$ ,

such that

- the isomorphisms respect the 2-functorial structure of  $h_X$  and  $k_X$ ,
- all 1-cells in  $\mathcal{V}$  are sent to central 1-cells in  $Kl(T)$ ,
- all the data respects the associativity and units for  $\otimes$ , meaning that  $h_X h_{X'}$  is isomorphic to  $h_{X \otimes X'}$ , the dual for  $k$ , that  $h_X k_Y$  is isomorphic to  $k_Y h_X$ , and that  $h_I$  is isomorphic to the identity and dually for  $k_I$ , all with the isomorphisms on objects determined by those of  $\mathcal{V}$ , and
- the symmetry  $\mathbb{C}$  of  $\mathcal{V}$  extends to being 2-natural on  $Kl(T)$ .

## 6 Properties of dense $KZ$ -monads

Now we have the notion of pseudo-commutativity, we can investigate further properties of dense  $KZ$ -monads. For our first result, we need care with the precise statement for two reasons. In general, to give a strength to a monad on a symmetric monoidal closed  $\mathcal{V}$  is equivalent to giving a  $\mathcal{V}$ -enrichment of the monad. So every 2-monad on  $\mathbf{Cat}$  has a unique strength associated with it. If  $\mathcal{V}$  is symmetric monoidal closed, complete and cocomplete,  $\mathcal{V}\text{-Cat}$



acquires a symmetric monoidal closed structure as a 2-category [8], but a 2-monad on  $\mathcal{V}\text{-Cat}$  is not a  $\mathcal{V}\text{-Cat}$ -enriched monad on  $\mathcal{V}\text{-Cat}$ , so there is no reason in general to expect it to have a strength. In particular cases, it does have a strength, for instance if  $\mathcal{V}$  is the arrow category, and so  $\mathcal{V}\text{-Cat}$  is **Poset**. We resolve this by restricting attention to  $\mathcal{V}\text{-Cat}$ -enriched monads on  $\mathcal{V}\text{-Cat}$ . If  $\mathcal{V} = \mathbf{Cat}$ , this reduces to 2-monads on **Cat**.

**6.1 Example** Let  $F$  be any class of weights such that the  $F$ -cocompletion of any small  $\mathcal{V}$ -category is small. Then the induced 2-monad acquires an enrichment in  $\mathcal{V}\text{-Cat}$ . This follows by close inspection of left Kan extensions in enriched category theory [8].

Second, there is delicacy with coherence. In principle, we should like to prove that dense  $KZ$ -monads are always commutative. But we cannot prove precisely that, but can only prove the result up to canonical isomorphism. Here, we need the definition of pseudo-commutativity we developed in Section 5. By the usual constructions involving change of universe, the definitions and results extend from small to large categories. So assuming that, we have

**6.2 Theorem** Every  $\mathcal{V}\text{-Cat}$ -monad  $T$  on  $\mathcal{V}\text{-Cat}$  whose underlying 2-monad is dense  $KZ$  has a pseudo-commutativity.

**Proof** This is most easily proved by use of Theorem 4.2. The two  $\mathcal{V}$ -functors from  $T\mathbb{C} \otimes T\mathbb{D}$  to  $T(\mathbb{C} \otimes \mathbb{D})$  send  $(f, g)$  to  $\text{colim}(f, \text{colim}(g, Y))$  and  $\text{colim}(g, \text{colim}(f, Y))$ . Since colimits commute with colimits, and because comparison maps are unique, we have a pseudo-commutativity. ■

**6.3 Corollary** If  $T$  is a  $\mathcal{V}\text{-Cat}$ -monad on  $\mathcal{V}\text{-Cat}$  whose underlying 2-monad is dense  $KZ$ , then  $Kl(T)$  is equivalent to a symmetric *Gray*-monoid with monoidal operation given by  $T(C \otimes D)$ .

A detailed description of the structure of  $Kl(T)$  rather than that of a 2-category equivalent to  $Kl(T)$  is given by Theorem 5.8.

We now consider constructions of the form  $T\mathbb{C} \times T\mathbb{D}$ . Here, we can make a slightly stronger statement by an axiomatic proof than we can by using an explicit construction of an equivalent version of  $T$ . The latter would yield an equivalence of  $\mathcal{V}$ -categories, whereas axiomatically we can obtain an

isomorphism, avoiding unpleasant coherence issues. We need to assume  $\mathcal{V}$  cartesian closed here, as we use constant  $\mathcal{V}$ -functors.

**6.4 Theorem** Let  $\mathcal{V}$  be cartesian closed and let  $T$  be a dense  $KZ$ -monad on  $\mathcal{V}\text{-Cat}$ . Suppose every  $T\mathbb{C}$  has and every  $TH$  and  $\mu_{\mathbb{C}}$  strictly preserve finite coproducts. Then  $T\mathbb{C} \times T\mathbb{D}$  is isomorphic to  $T(\mathbb{C} + \mathbb{D})$  2-naturally in  $\mathbb{C}$  and  $\mathbb{D}$  and coherently with respect to the associative, commutative, and unit structures of binary product and coproduct.

**Proof** First observe that every  $\mathcal{V}$ -functor in  $Kl(T)$  strictly preserves finite coproducts. Using the universal property of the Kleisli construction, and using the universal properties of products and coproducts, and in order to define a  $\mathcal{V}$ -functor  $H : T(\mathbb{C} + \mathbb{D}) \rightarrow T\mathbb{C} \times T\mathbb{D}$ , we give  $\mathcal{V}$ -functors from  $\mathbb{C}$  to each of  $T\mathbb{C}$  and  $T\mathbb{D}$  and from  $\mathbb{D}$  to each of  $T\mathbb{C}$  and  $T\mathbb{D}$ . We define them by  $\eta_{\mathbb{C}}$  and the constant at the initial object of  $T\mathbb{D}$ , and by duality.

We define  $K : T\mathbb{C} \times T\mathbb{D} \rightarrow T(\mathbb{C} + \mathbb{D})$  by sending  $(X, Y)$  to  $(Ti_0)X + (Ti_1)Y$ , where  $i_0 : \mathbb{C} \rightarrow \mathbb{C} + \mathbb{D}$  and  $i_1 : \mathbb{D} \rightarrow \mathbb{C} + \mathbb{D}$  are the left and right coprojections respectively.

We must show that  $H$  and  $K$  are mutually inverse. They are obviously 2-natural in  $\mathbb{C}$  and  $\mathbb{D}$ . To see that  $KH = id_{T(\mathbb{C} + \mathbb{D})}$ , first see that  $KH\eta_{\mathbb{C} + \mathbb{D}} = \eta_{\mathbb{C} + \mathbb{D}}$ , which may be checked on each component. That follows routinely since  $Ti_0$  strictly preserves the initial object, and dually. Now, using the explicit coherence assumption of the theorem and some routine manipulation of diagrams, the commutativity extends to  $T(\mathbb{C} + \mathbb{D})$ .

To see that  $HK = id_{T\mathbb{C} \times T\mathbb{D}}$ , we must verify that  $\pi_0 HK = \pi_0$  and  $\pi_1 HK = \pi_1$ , where  $\pi_0$  and  $\pi_1$  are the first and second projections from  $T\mathbb{C} \times T\mathbb{D}$  respectively. By definition,  $\pi_0 H$  and  $\pi_1 H$  both lie in  $Kl(T)$ , so preserve finite coproducts strictly. Restricting our attention to  $\pi_0$ , the other case being dual, it suffices to show that  $(\pi_0 H \times \pi_0 H)(Ti_0 \times Ti_1)$  sends  $(X, Y)$  to  $(X, 0)$ , where  $0$  is the initial object of  $T\mathbb{C}$ . Again, this amounts to two commutativities.

For the first, observe that  $\pi_0 H = \mu_{\mathbb{C}} T(\eta_{\mathbb{C}}, 0)$ , so precomposing with  $Ti_0$  yields the identity since  $(\eta_{\mathbb{C}}, 0)i_0 = \eta_{\mathbb{C}}$  and by one of the monadic unit laws, giving the desired commutativity.

For the second, by a similar calculation, it suffices to show that the lifting of the constant  $\mathcal{V}$ -functor  $0 : \mathbb{D} \rightarrow T\mathbb{C}$  to  $T\mathbb{D}$  is the constant  $\mathcal{V}$ -functor at the initial object  $0$  of  $T\mathbb{C}$ . But by Theorem 3.7, the lifting is given by the left Kan extension of  $0 : \mathbb{D} \rightarrow T\mathbb{C}$  along  $\eta_{\mathbb{C}} : \mathbb{C} \rightarrow T\mathbb{C}$ ; and one can check

by calculation that that is necessarily the constant at 0. ■

**6.5 Corollary** Under the hypotheses of the Theorem,  $Kl(T)$  has finite products and finite coproducts, both given by  $T\mathbb{C} \times T\mathbb{D}$ .

**Proof** Observe by construction that the projections from  $T(\mathbb{C} \times \mathbb{D})$  to each of  $T\mathbb{C}$  and  $T\mathbb{D}$  are in  $Kl(T)$  and the compositions used in the proof of the Theorem agree with those in  $Kl(T)$ . ■

## References

- [1] M.H. Albert and G.M. Kelly, The closure of a class of colimits, *J. Pure Appl. Algebra* 51 (1988) 1–17.
- [2] R. Blackwell, G.M. Kelly, and A.J. Power, Two-dimensional monad theory, *J. Pure Appl. Algebra* 59 (1989) 1–41.
- [3] G. L. Cattani, A.J. Power, and G. Winskel, A categorical axiomatics for bisimulation, In *Proceedings of CONCUR '98*, Lecture Notes in Computer Science 1466 (1998) 581–596.
- [4] R. Gordon and A.J. Power, Gabriel Ulmer duality for categories enriched in bicategories, *J. Pure Appl. Algebra* (to appear).
- [5] R. Gordon, A.J. Power, and Ross Street, Coherence for tricategories, *Memoirs Amer. Math. Soc.* 558 (1995).
- [6] A. Joyal, M. Nielsen, and G. Winskel, Bisimulation from open maps, *Information and Computation* 127 (1996) 164–185.
- [7] G.M. Kelly, Coherence conditions for lax algebras and for distributive laws, *Proc. Sydney Category Theory Seminar 1972/73*, Lecture Notes in Math. 420, Springer (1974) 281–375.
- [8] G.M. Kelly, Basic concepts of enriched category theory, *Notes London Math. Soc Lecture Series* 64, Cambridge University Press (1982).
- [9] G.M. Kelly and R. Street, Review of the elements of 2-categories, *Proc. Sydney Category Theory Seminar 1972/73*, Lecture Notes in Math. 420, Springer (1974) 75–103.

- [10] A. Kock, Closed categories generated by commutative monads, *J. Austral. Math. Soc.* 12 (1971) 405–424.
- [11] A. Kock, Monads for which structures are adjoints to units, *J. Pure Appl. Algebra* 104 (1995) 41–59.
- [12] R. Milner, *Communication and concurrency*, Prentice Hall (1989).
- [13] A.J. Power and E.P. Robinson, Premonoidal categories and notions of computation, *Math. Structures in Computer Science* 7 (1997) 453–468.

## Recent BRICS Report Series Publications

- RS-98-21 John Power, Gian Luca Cattani, and Glynn Winskel. *A Representation Result for Free Cocompletions*. September 1998. 16 pp.
- RS-98-20 Søren Riis and Meera Sitharam. *Uniformly Generated Submodules of Permutation Modules*. September 1998. 35 pp.
- RS-98-19 Søren Riis and Meera Sitharam. *Generating Hard Tautologies Using Predicate Logic and the Symmetric Group*. September 1998. 13 pp.
- RS-98-18 Ulrich Kohlenbach. *Things that can and things that can't be done in PRA*. September 1998. 24 pp.
- RS-98-17 Roberto Bruni, José Meseguer, Ugo Montanari, and Vladimiro Sassone. *A Comparison of Petri Net Semantics under the Collective Token Philosophy*. September 1998. 20 pp. To appear in *4th Asian Computing Science Conference, ASIAN '98 Proceedings, LNCS, 1998*.
- RS-98-16 Stephen Alstrup, Thore Husfeldt, and Theis Rauhe. *Marked Ancestor Problems*. September 1998.
- RS-98-15 Jung-taek Kim, Kwangkeun Yi, and Olivier Danvy. *Assessing the Overhead of ML Exceptions by Selective CPS Transformation*. September 1998. 31 pp. To appear in the proceedings of the *1998 ACM SIGPLAN Workshop on ML*, Baltimore, Maryland, September 26, 1998.
- RS-98-14 Sandeep Sen. *The Hardness of Speeding-up Knapsack*. August 1998. 6 pp.
- RS-98-13 Olivier Danvy and Morten Rhiger. *Compiling Actions by Partial Evaluation, Revisited*. June 1998. 25 pp.
- RS-98-12 Olivier Danvy. *Functional Unparsing*. May 1998. 7 pp. This report supersedes the earlier report BRICS RS-98-5. Extended version of an article to appear in *Journal of Functional Programming*.
- RS-98-11 Gudmund Skovbjerg Frandsen, Johan P. Hansen, and Peter Bro Miltersen. *Lower Bounds for Dynamic Algebraic Problems*. May 1998. 30 pp.