



Basic Research in Computer Science

BRICS RS-94-3

U. H. Engberg & G. Winskel: Linear Logic on Petri Nets

Linear Logic on Petri Nets

Uffe H. Engberg
Glynn Winskel

BRICS Report Series

RS-94-3

ISSN 0909-0878

February 1994

**Copyright © 1994, BRICS, Department of Computer Science
University of Aarhus. All rights reserved.**

**Reproduction of all or part of this work
is permitted for educational or research use
on condition that this copyright notice is
included in any copy.**

**See back inner page for a list of recent publications in the BRICS
Report Series. Copies may be obtained by contacting:**

**BRICS
Department of Computer Science
University of Aarhus
Ny Munkegade, building 540
DK - 8000 Aarhus C
Denmark
Telephone: +45 8942 3360
Telefax: +45 8942 3255
Internet: BRICS@daimi.aau.dk**

Linear Logic on Petri Nets

Uffe H. Engberg

Glynn Winskel

BRICS*

Computer Science Department[†]

Aarhus University

Ny Munkegade

DK-8000 Aarhus C, Denmark

ABSTRACT. This article shows how individual Petri nets form models of Girard's intuitionistic linear logic. It explores questions of expressiveness and completeness of linear logic with respect to this interpretation. An aim is to use Petri nets to give an understanding of linear logic and give some appraisal of the value of linear logic as a specification logic for Petri nets. This article might serve as a tutorial, providing one in-road into Girard's linear logic via Petri nets. With this in mind we have added several exercises and their solutions. We have made no attempt to be exhaustive in our treatment, dedicating our treatment to one semantics of intuitionistic linear logic.

Completeness is shown for several versions of Girard's linear logic with respect to Petri nets as the class of models. The strongest logic considered is intuitionistic linear logic, with \otimes , \multimap , $\&$, \oplus and the exponential ! ("of course"), and forms of quantification. This logic is shown sound and complete with respect to *atomic nets* (these include nets in which every transition leads to a nonempty multiset of places). The logic is remarkably expressive, enabling descriptions of the kinds of properties one might wish to show of nets; in particular, negative properties, asserting the impossibility of an assertion, can also be expressed. A start is made on decidability issues.

KEYWORDS. Linear logic, Petri nets.

Contents

1	Introduction	2
2	Intuitionistic Linear Logic	5
3	Proof Rules	5
4	Quantale Interpretation	8
5	Petri Nets	10

*Basic Research in Computer Science, Centre of the Danish National Research Foundation.

[†]e-mail address: {engberg,gwinskel}@daimi.aau.dk, fax: +45 89 42 32 55

7	Petri-Net Interpretation	13
8	Expressing Properties of Nets	16
9	The “Of Course” Operator	19
10	Linear Negation	20
11	Recursion	23
12	Equivalences on Nets	24
13	Elementary Completeness Results	26
14	Quantification and Atomic Nets	33
15	Decidability Issues	47
16	Conclusion	50
A	Answers to exercises	51

1 Introduction

Girard’s linear logic arose in part from Girard’s insights into particular categorical models for intuitionistic logic. In giving a domain-theoretic semantics to his System F (the polymorphic λ -calculus) [Gir86], Girard re-invented the stable domain theory of Berry based on the category dI-domains and stable functions [Ber78], though for rather special domains, the coherence spaces. It turns out that Girard’s construction for modelling polymorphic types can be carried through, with some modifications, in the more standard domain theory based on the category of Scott domains and continuous functions (see [CGW89]). But Girard’s use of the category of coherence spaces and stable functions led to a significant discovery, that of classical linear logic. Girard recognised that the category of coherence spaces with stable functions contained a reflective subcategory in which morphisms were “linear”; the adjunction explained how the larger category could be derived from the linear one, and through the propositions-as-types reading of logic it gave an explanation of intuitionistic logic in terms of a more primitive and new classical linear logic. Again, it turns out a similar observation could have been made were Girard to have used Scott’s domain theory, though the linear logic would then have had less structure (that of *intuitionistic* rather than *classical* linear logic). This is a thumbnail sketch of the discovery of linear logic, and indicates why it is

Right from the start [Gir87], the categorical model of coherence spaces played a prominent role in the presentation of linear logic, and since other categorical models, arguably of a more informative character, have appeared; notable are those based on game semantics, initiated by Blass’s work (see [Bla92, AJ92]). Not everyone is happy with a categorical model as the explanation of a new logic, and in his pioneering paper [Gir87], Girard also gave a “phase–space” semantics for linear logic. There have since been investigations of several other kinds of structures as models of linear logic, quantales (e.g.[AV88]), generalised Kripke structures [AD93], and Petri nets—the main topic here.

Girard’s linear logic has sparked off a great deal of interest in how it might be useful in the theory of parallelism, not least because of Girard’s initial claims for it [Gir87]. Linear logic has been described as a “resource conscious” logic by Martí-Oliet and Meseguer [MOM89]; in its proofs occurrences of propositions cannot be used more than once or disappear unless they are explicitly created or used up by the rules of inference. People were not long in spotting a relationship with Petri nets where there are similar ideas. Places in a Petri net hold to certain nonnegative multiplicities forming a multiset of places, traditionally called a marking; as transitions occur, multiplicities of places are consumed and produced in accord with a dynamic behaviour of nets, formalised in the so-called “token game”. Independently, Gunter and Gehlot [GG89a], Asperti [Asp87] and Brown [Bro89] showed that places are like atomic propositions in linear logic and transitions like proof rules. In [GG89a] and [Asp87], the fine grain of linear logic proofs for the \otimes -fragment of linear logic is related to the token game in Petri nets. Essentially it is shown how a proof of $A \vdash B$ in linear logic, where A and B are built from place names just with \otimes , corresponds to a play of the token game taking the marking corresponding to A to that corresponding to B .

The work of Martí-Oliet and Meseguer [MOM89] extends that of [GG89a] and [Asp87]. It is known that certain kinds of “linear” categories are models for linear logic. The work [MOM89] essentially proceeds by letting Petri nets freely generate a linear category and then interpreting linear logic in that setting. One problem with their approach is its consequence that if a net satisfies an proposition of linear logic then so does any augmentation of the net, obtained by adding transitions arbitrarily. This considerably weakens the case for regarding linear logic as a specification logic with respect to *their* notion of satisfaction.¹ Nor at this exploratory stage, when we are trying to understand what use linear logic might be on Petri nets, is it so clear, that a linear category with lots of proof terms as morphisms has any advantage over a more accessible partial order semantics, of the kind presented here.

¹No one would take seriously a program logic with the property that if a program satisfied an assertion then so did the program with GOTO statements inserted arbitrarily.

of linear logic had arisen before in the form of quantales. Indeed Girard’s phase semantics in [Gir87] for linear logic uses free quantales. Abramsky and Vickers [AV88] approached quantales from a computer science viewpoint, the hope being that it would lead to a “linear process logic”. Yetter [Yet] and Rosenthal [Ros89] looked at quantales and linear logic more from the perspective of pure mathematics—how to represent them and their relationship with other bits of mathematics.

We point out a straightforward way in which a Petri net induces a quantale and so becomes a model for intuitionistic linear logic. The model is for all of intuitionistic linear logic. A prime feature is its accessibility; this is important with the new logic, where working out what you can and cannot say is tricky. That it generalises the work of [GG89a] and [Asp87] is clear. The paper provides evidence that intuitionistic linear logic, with the right notion of satisfaction, can be a reasonably expressive specification logic for parallel processes. It also throws some light on the use of classical linear logic; via a construction, observed by Abramsky and Vickers [AV88], generalising that of Girard’s for phase semantics, a quantale can easily be turned into a model for classical linear logic, although unfortunately when taken to nets, the resulting semantics of classical linear logic is often trivial.

Our strongest completeness result is for the full logic described in [GL87, Laf88], viz. it includes

$$\otimes, \multimap, \oplus, \&, \text{ and } !$$

though at a cost, to the purity of the linear logic, of adding quantification over markings and axioms special to the net semantics. For this strongest completeness result, a slight restriction is also made to the Petri nets considered as models; they should be *atomic* (see definition 36), but fortunately this restriction is one generally met, and even often enforced, in working with Petri nets. The step of considering only atomic nets as models has two important pay-offs: one is that the exponential $!A$ becomes definable as $A \& \mathbf{1}$, where $\mathbf{1}$ is the unit of \otimes ; the other is that we can say internally, within the logic, that an assertion is *not* satisfied—the possibility of asserting such negative properties boosts the logic’s expressive power considerably. We can achieve completeness for more modest fragments of the logic without extra axioms and with respect to the entire class of nets as models (see section 13).

The work here (filling out that of [EW90, EW93]) contrasts with other approaches to linear logic on Petri nets in that they either apply only to much smaller fragments of the logic such as the \otimes -fragment (cf. [GG89b]), or use the transitions of a Petri net to freely generate a linear-logic theory (cf. [MOM91]), in which case the logic becomes rather inexpressive, and in particular cannot capture negative

2 Intuitionistic Linear Logic

In the Gentzen sequent calculus for intuitionistic logic a sequent, $A_1, \dots, A_n \vdash A$, is written to mean that the formula A is deducible from the assumption formulae A_1, \dots, A_n (we shall use Γ as an abbreviation for a (possibly empty) sequence of assumption formulae). The calculus has the two structural rules

$$\frac{\Gamma \vdash B}{\Gamma, A \vdash B} \text{(thinning)} \quad \frac{\Gamma, A, A \vdash B}{\Gamma, A \vdash B} \text{(contraction)} \quad (1)$$

for adding an assumption and removal of a duplicate. In the presence of these rules the following two right introduction rules for conjunction

$$\frac{\Gamma \vdash A \quad \Delta \vdash B}{\Gamma, \Delta \vdash A \wedge B} \quad \frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B} \quad (2)$$

become interderivable in the sense that the first rule can be derived from the second by thinning, and the second from the first by contraction.

Now, as remarked in [AV88], if a proof is to be seen as representing the *process* of proving, then these structural rules become far from self-evident. Indeed, in intuitionistic linear logic these rules are dropped and the rules of (2) are no longer interderivable. Without them, propositions cannot be introduced arbitrarily into a list of assumptions and nor can a duplication in the assumptions be removed. It is in this sense that linear logic is a “resource conscious” logic. As a consequence of dropping (1) the two rules, (2), become two separate right introduction rules for two fundamentally different forms of conjunction; they are written as \otimes and $\&$ respectively. These new connectives are part of Girard’s intuitionistic linear logic, which we shall now present.

3 Proof Rules

The connectives of intuitionistic linear logic are:

- \otimes tensor, with unit **1**, called one,
- $\&$ conjunction, with unit **T**, called true,
- \oplus disjunction, with unit **F**, called false.

We take as the definition of intuitionistic linear logic the proof rules presented in [GL87, Laf88]:

$$\frac{}{A \vdash A} \text{(identity)} \quad \frac{\Gamma \vdash A \quad \Delta, A \vdash B}{\Gamma, \Delta \vdash B} \text{(cut)} \quad \frac{\Gamma, A, B, \Delta \vdash C}{\Gamma, B, A, \Delta \vdash C} \text{(exchange)}$$

Logical rules

$$\frac{\Gamma \vdash A \quad \Delta \vdash B}{\Gamma, \Delta \vdash A \otimes B} (\vdash \otimes) \quad \frac{\Gamma, A, B \vdash C}{\Gamma, A \otimes B \vdash C} (\otimes \vdash) \quad \frac{\Gamma \vdash A}{\Gamma, \mathbf{1} \vdash A} \quad \frac{}{\vdash \mathbf{1}}$$

$$\frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \& B} (\vdash \&) \quad \frac{\Gamma, A \vdash C}{\Gamma, A \& B \vdash C} (l\&\vdash) \quad \frac{\Gamma, B \vdash C}{\Gamma, A \& B \vdash C} (r\&\vdash) \quad \frac{}{\Gamma \vdash \mathbf{T}}$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash A \oplus B} (\vdash \oplus l) \quad \frac{\Gamma \vdash B}{\Gamma \vdash A \oplus B} (\vdash \oplus r) \quad \frac{\Gamma, A \vdash C \quad \Gamma, B \vdash C}{\Gamma, A \oplus B \vdash C} (\oplus \vdash) \quad \frac{}{\Gamma, \mathbf{F} \vdash A}$$

$$\frac{\Gamma, A \vdash B}{\Gamma \vdash A \multimap B} (\vdash \multimap) \quad \frac{\Gamma \vdash A \quad \Delta, B \vdash C}{\Gamma, \Delta, A \multimap B \vdash C} (\multimap \vdash)$$

As an example of a proof, we can derive the rule $\frac{\Gamma \vdash A \multimap B}{\Gamma, A \vdash B}$ from $(\multimap \vdash)$ by

$$\frac{\frac{\frac{}{A \vdash A} \quad \frac{}{B \vdash B}}{\Gamma \vdash A \multimap B} \quad \frac{}{A, A \multimap B \vdash B}}{\Gamma, A \vdash B}}{.}$$

The absence of the rules for thinning and contraction is compensated, to some extent, by the addition of the logical operator “of course”. In [GL87, Laf88] this operator is presented with the following proof rules (stronger than those in [Gir87]):

“Of course” rules

$$\frac{}{\mathbf{!}A \vdash A} \quad \frac{}{\mathbf{!}A \vdash \mathbf{1}} \quad \frac{}{\mathbf{!}A \vdash \mathbf{!}A \otimes \mathbf{!}A} \quad (3)$$

$$\frac{B \vdash A \quad B \vdash \mathbf{1} \quad B \vdash B \otimes B}{B \vdash \mathbf{!}A} \quad (4)$$

Given a proposition A , the assertion of $\mathbf{!}A$ has the possibility of being instantiated by the proposition A , the unit $\mathbf{1}$ or $\mathbf{!}A \otimes \mathbf{!}A$, and thus of arbitrarily many assertions of $\mathbf{!}A$.

How this operator compensates for the absence of the two structural rules can be seen from the (weaker) derived rules Girard originally presented in [Gir87]

$$\frac{\Gamma, A \vdash B}{\Gamma, \mathbf{!}A \vdash B} \quad \frac{\Gamma \vdash B}{\Gamma, \mathbf{!}A \vdash B} \quad \frac{\Gamma, \mathbf{!}A, \mathbf{!}A \vdash B}{\Gamma, \mathbf{!}A \vdash B} \quad (5)$$

and

$$\overline{!\Gamma \vdash !B} \quad (6)$$

where $!\Gamma$ is a shorthand for $!A_1, \dots, !A_n$ when $\Gamma = A_1, \dots, A_n$.

Once a logical constant, \perp , denoting linear-absurdity is fixed on, linear negation (not to be mistaken for intuitionistic negation) is derivable:

$$A^\perp = A \multimap \perp .$$

We end the section by showing that the original “of course” rules indeed are derivable.

(5) is derived from (3) as follows

$$\frac{\overline{!A \vdash A} \quad \Gamma, A \vdash B}{\Gamma, !A \vdash A} \quad \frac{\overline{!A \vdash \mathbf{1}} \quad \frac{\Gamma \vdash B}{\Gamma, \mathbf{1} \vdash B}}{\Gamma, !A \vdash B} \quad \frac{\overline{!A \vdash !A \otimes !A} \quad \frac{\Gamma, !A, !A \vdash B}{\Gamma, !A \otimes !A \vdash B}}{\Gamma, !A \vdash B}$$

leaving out use of the exchange rule. (6) is however not so directly derivable from (4). At first we derive

$$\frac{!A \vdash B}{!A \vdash !B} \quad \text{and} \quad \frac{A \vdash B}{!A \vdash !B} \quad (7)$$

from (4) by

$$\frac{!A \vdash B \quad \overline{!A \vdash \mathbf{1}} \quad \overline{!A \vdash !A \otimes !A}}{!A \vdash !B} \quad \text{and} \quad \frac{\overline{!A \vdash A} \quad A \vdash B}{!A \vdash !B} .$$

We are then in a position to derive

$$!(A \& B) \dashv\vdash !A \otimes !B$$

where $A \dashv\vdash B$ means $A \vdash B$ and $B \vdash A$ are provable. Using (4) the first proof is

$$\frac{\frac{\overline{!A \vdash A} \quad \overline{!B \vdash \mathbf{1}}}{!A \otimes !B \vdash A \otimes \mathbf{1}} \quad \frac{\overline{!A \vdash \mathbf{1}} \quad \overline{!B \vdash B}}{!A \otimes !B \vdash \mathbf{1} \otimes B} \quad \frac{\overline{!A \vdash \mathbf{1}} \quad \overline{!B \vdash \mathbf{1}}}{!A \otimes !B \vdash \mathbf{1} \otimes \mathbf{1}} \quad \frac{\overline{!A \vdash !A \otimes !A} \quad \overline{!B \vdash !B \otimes !B}}{!A \otimes !B \vdash (!A \otimes !A) \otimes (!B \otimes !B)}}{\frac{!A \otimes !B \vdash A \quad !A \otimes !B \vdash B}{!A \otimes !B \vdash A \& B} \quad \frac{!A \otimes !B \vdash \mathbf{1} \otimes \mathbf{1} \quad !A \otimes !B \vdash (!A \otimes !B) \otimes (!A \otimes !B)}}{!A \otimes !B \vdash !(A \& B)}}$$

and the other proof follows from the second rule of (7) by:

$$\frac{\frac{\overline{A \vdash A} \quad \overline{B \vdash B}}{A \& B \vdash A} \quad \frac{\overline{B \vdash B}}{A \& B \vdash B}}{!(A \& B) \vdash !A \quad !(A \& B) \vdash !B} \quad \frac{!(A \& B) \vdash !A \quad !(A \& B) \vdash !B}{!(A \& B), !(A \& B) \vdash !A \otimes !B}}{\frac{!(A \& B) \vdash !(A \& B) \otimes !(A \& B) \quad !(A \& B) \otimes !(A \& B) \vdash !A \otimes !B}{!(A \& B) \vdash !A \otimes !B}} .$$

$$\frac{\Gamma \vdash B}{\otimes \Gamma \vdash B} \quad \frac{\otimes \Gamma \vdash B}{\Gamma \vdash B}$$

we get

$$!(\& \Gamma) \dashv\vdash \otimes !\Gamma$$

where for $\Gamma = A_1, \dots, A_n$, $\otimes \Gamma$ means $A_1 \otimes \dots \otimes A_n$ and similarly for $\& \Gamma$. With this we then obtain (6) from the first rule of (7).

Exercise 1 Show the following:

$$(i) \frac{\Gamma \vdash A \& B}{\Gamma \vdash A} \quad \text{and} \quad \frac{\Gamma, A \otimes B \vdash C}{\Gamma, A, B \vdash C}$$

$$(ii) (A \oplus B) \multimap C \vdash (A \multimap C) \& (B \multimap C)$$

$$(iii) A \otimes (B \oplus C) \dashv\vdash (A \otimes B) \oplus (A \otimes C) \quad \square$$

4 Quantale Interpretation

We have just seen the proof rules of intuitionistic linear logic. What are its models? As recognised by several people [AV88, Yet, Ros89, Sam], quantales² provide an algebraic semantics for intuitionistic linear logic. Quantales are to intuitionistic linear logic as complete Heyting algebras are to intuitionistic logic. A quantale is a commutative monoid on a complete join semilattice. Spelled out:

Definition 2 *A quantale Q is a complete join semilattice (i.e. a partial order with an operation forming joins of arbitrary sets) together with an associative, commutative, binary operation \otimes and constant $\mathbf{1}$ such that*

$$q \otimes \mathbf{1} = q$$

$$q \otimes \bigvee P = \bigvee \{q \otimes p \mid p \in P\} \quad \square$$

Entailment is interpreted as the order relation, \leq , on the underlying lattice of a quantale. The logical operation, \otimes , is interpreted by the corresponding binary operation in the quantale and the logical constant $\mathbf{1}$ is interpreted as $\mathbf{1}$ in the quantale. The disjunction, \oplus , of linear logic is understood as binary join and the conjunction, $\&$, as binary meet. The logical constants \mathbf{T} and \mathbf{F} are interpreted as the top and bottom element respectively of the complete lattice. Linear implication is a derived operation

$$p \multimap q \stackrel{def}{=} \bigvee \{r \mid r \otimes p \leq q\}$$

²As originally defined, quantales need not be commutative and should satisfy the idempotency law $q \otimes q = q$. We shall take quantales to be commutative and relax the idempotency law.

a complete Heyting algebra, but this time w.r.t. \otimes in place of \wedge . The definition of linear implication ensures the adjunction:

$$r \otimes p \leq q \quad \text{iff} \quad r \leq p \multimap q .$$

With respect to a quantale, and interpretations of the atomic propositions as elements of a quantale, we can inductively associate a proposition A in linear logic with its denotation as a quantale element $\llbracket A \rrbracket$. An entailment

$$A_1, \dots, A_n \Vdash A$$

holds in the quantale iff

$$\llbracket A_1 \rrbracket \otimes \dots \otimes \llbracket A_n \rrbracket \leq \llbracket A \rrbracket .$$

The special case where $n = 0$ is allowed, in which case the situation amounts to

$$\Vdash A \quad \text{iff} \quad \mathbf{1} \leq \llbracket A \rrbracket .$$

It is a routine matter to check that each rule is sound with respect to this interpretation. For example the right and left introduction rules for disjunction, \oplus , and conjunction, $\&$, express that they are the join and meet with respect to entailment while the rules for linear implication express the adjunction which characterise it (with the help of the little proof given as an example in section 2). In this way it can be seen that, with respect to a quantale:

Theorem 3 *If $\vdash A$ then $\Vdash A$.*

We have so far ignored the treatment of $!A$. The rules of (3) for $!A$ are interderivable with the following single rule:

$$\overline{!A \vdash \mathbf{1} \& A \& (!A \otimes !A)}$$

So, as an interpretation of $!q$, for an element q of a quantale, we require an element x such that

$$x \leq \mathbf{1} \& q \& (x \otimes x) . \tag{8}$$

This will not in general characterise a unique value of the quantale; for instance taking x to be the bottom element of the lattice will always do. However from (4) it follows that any x satisfying (8) should be below $!q$, and hence $!q$ should be the greatest postfixing point, and so fixed point, of

$$x \mapsto \mathbf{1} \& q \& (x \otimes x)$$

in the complete lattice given together with the quantale. Such a solution ensures the soundness of the proof rules extended by those for $!A$.

quantale is determined by a choice for the denotation of \perp . Then A^\perp is understood as $A \multimap \perp$. In this abstract set-up, for a general quantale we cannot hope to say much more. However, in a moment, when considering quantales got from a Petri net, a special element will be proposed as being useful. We can observe, following Abramsky and Vickers [AV88], that from a choice for \perp in a quantale, we can obtain a model of classical linear logic. The key feature of the classical version is that linear negation is an involution. With respect to an element \perp of a quantale \mathcal{Q} , write q^\perp for $q \multimap \perp$. We cannot expect $q = q^{\perp\perp}$ to hold for a general quantale. However, the image of the map $(-)^{\perp\perp} : \mathcal{Q} \rightarrow \mathcal{Q}$ is a subquantale of \mathcal{Q} on which $(-)^{\perp}$ is involutive, so forming a model of classical linear logic.

Exercise 4

1. Show that \otimes is monotonic (recall $p \leq q \Leftrightarrow p \vee q = q$).
2. Prove the adjunction $r \otimes p \leq q$ iff $r \leq p \multimap q$. □

5 Petri Nets

Petri nets are a model of processes (or systems) in terms of types of resources, represented by *places* which can hold to arbitrary nonnegative multiplicity, and how those resources are consumed or produced by actions, represented by *transitions*. They are described using the notation of multisets.

A *multiset* over a set P is a function, $M : P \rightarrow \mathbb{N}$. We shall henceforth only be concerned with *finite* multisets, i.e. $\{a \in P \mid M(a) \neq 0\}$ finite. With addition, $+$, of multisets defined by $(M + M')(a) = M(a) + M'(a)$ for all $a \in P$, multisets over P form a (free) commutative monoid with $\underline{0}$ ($\forall a \in P. \underline{0}(a) = 0$), the empty multiset, as unit. An element a of P can be turned into the singleton multiset with multiplicity 1 at a and 0 elsewhere. Observe that multisets are partially ordered by $M \leq M'$ iff $\forall a \in P. M(a) \leq M'(a)$. For $n \in \mathbb{N}$ and M a multiset, the scalar multiplication nM is defined by $(nM)(a) = n \cdot M(a)$.

We take a Petri net N to consist of $(P, T, \bullet(-), (-)^\bullet)$, where P , a set of *places*, and T , a set of *transitions*, are accompanied by maps $\bullet(-), (-)^\bullet$ on transitions T which for each $t \in T$ give a multiset of P , called the pre- and post (multi)set of t respectively. For the moment there are none of the usual restrictions on the net, such as absence of isolated elements, and in particular transitions with empty pre sets and/or post sets will be allowed. And we are actually considering nets with unconstrained capacity.

tribution of resources, formalized in the definition of a marking. A *marking* of N will simply be a finite multiset over P . We use \mathcal{M} to denote the *set of markings* of the net, understood from the context. Sometimes nets are associated with an initial marking M_0 . The behaviour of a net is expressed by saying how markings change as transitions occur (or fire). For markings, M, M' , and a transition $t \in T$, $M [t\rangle M'$ stands for t fires from M to M' ; that is the firing relation $[t\rangle$ is given by

$$M [t\rangle M' \quad \text{iff} \quad \exists M'' \in \mathcal{M}. M = M'' + \bullet t \text{ and } t\bullet + M'' = M' \quad .$$

So t is enabled at M if there is an $M' \in \mathcal{M}$ such that $M [t\rangle M'$. We shall write $M \rightarrow M'$ for the reachability relation, the reflexive and transitive closure of the firing relations.³

$$M \rightarrow M' \quad \text{iff} \quad \begin{aligned} &\exists t_1, \dots, t_n \in T, M_1, \dots, M_n \in \mathcal{M}, n \geq 0. \\ &M [t_1\rangle M_1 \cdots [t_n\rangle M_n = M' \end{aligned}$$

The markings, $[M\rangle$, forwards reachable from M form the set $\{M' \in \mathcal{M} \mid M \rightarrow M'\}$. We shall use $\downarrow M$ to denote the set of markings which can reach M . We will generally call this set the downwards closure of M . It is defined by $\downarrow M = \{M' \in \mathcal{M} \mid M' \rightarrow M\}$. We shall also use this notation, extended pointwise, on sets of markings H , taking e.g. $\downarrow H = \{M' \mid \exists M \in H. M' \rightarrow M\}$.

Petri nets can be presented by using the well-known graphical notation, which will be used throughout this paper. Places are represented by circles, transitions as squares, and arcs of appropriate multiplicities used to indicate the pre and post sets. The formal definitions can then be brought to life in the so called “token game” where markings are visualised as consisting of a distribution of tokens over places; the number of tokens residing on a place expresses the multiplicity to which it holds according to the marking. The tokens are consumed and produced as transitions occur. A basic reference for Petri nets is [Rei85].

Since we are dealing with place/transition nets without capacities we have the following proposition which is crucial for the net semantics of linear logic in the sequel.

Proposition 5 *For any $M, M', M'' \in \mathcal{M}$ we have:*

$$M \rightarrow M' \quad \text{implies} \quad M + M'' \rightarrow M' + M'' \quad .$$

³Because we shall be chiefly concerned with the reachability relation between markings, we can get by with single steps in the firing relation, and ignore the variant in which transitions can fire concurrently.

From a given net N , and associated markings and reachability relation \mathcal{M}, \rightarrow a quantale, \mathcal{Q} , can be constructed. Its elements are to be the downwards closed subsets of markings; a subset H of \mathcal{M} is downwards closed iff it satisfies

$$M' \rightarrow M \text{ and } M \in H \quad \text{implies} \quad M' \in H .$$


The set \mathcal{Q} is ordered by inclusion, \subseteq . With respect to inclusion, it is easy to see that union and intersection form the join and meet operations. The binary operation \otimes has as unit element: $\mathbf{1} = \downarrow \underline{0}$. For $p, q \in \mathcal{Q}$ the binary operation \otimes is defined by:

$$p \otimes q = \downarrow(p + q), \text{ where } p + q = \{M_p + M_q \mid M_p \in p, M_q \in q\} .$$

Thus

$$p \otimes q = \{M \mid \exists M_p \in p, M_q \in q. M \rightarrow M_p + M_q\} .$$

The downwards closure, \downarrow , in the definition of \otimes above is needed to make \otimes well-defined as can be seen from this net:



Clearly $\downarrow \underline{b} + \downarrow \underline{c} = \{M\} \neq \{\underline{a}, M\} = \downarrow \{M\}$, where $M = \underline{b} + \underline{c}$.

That \mathcal{Q} is indeed a commutative quantale follows from the commutativity of $+$ and the fact that for $p \in \mathcal{Q}$, and $Q \subseteq \mathcal{Q}$:

$$\begin{aligned} p \otimes \cup Q &= \downarrow(p + \cup Q) = \downarrow(\cup \{p + q \mid q \in Q\}) \\ &= \cup \{\downarrow(p + q) \mid q \in Q\} = \cup \{p \otimes q \mid q \in Q\} . \end{aligned}$$

Defining $p \multimap q = \cup \{r \mid r \otimes p \subseteq q\}$ we can simplify it to

$$p \multimap q = \{M \mid \forall M_p \in p. M + M_p \in q\} .$$

This is because

$$\begin{aligned} &M \in p \multimap q \\ \text{iff } &\exists r. M \in r, r \otimes p \subseteq q \\ \text{iff } &\downarrow M \otimes p \subseteq q, \quad \text{as } \downarrow M \in \mathcal{Q} \text{ and} \\ &M \in r \Rightarrow \downarrow M \subseteq r \\ \text{iff } &\forall M' \in \downarrow M, M_p \in p. M' + M_p \in q, \text{ as } q \text{ is downwards closed} \\ \text{iff } &\forall M_p \in p. M + M_p \in q \end{aligned}$$

where the “if” of the last equivalence follows from the downwards closure of q and $M' \in \downarrow M \Rightarrow M' \rightarrow M \Rightarrow M' + M_p \rightarrow M + M_p \Rightarrow M' + M_p \in \downarrow(M + M_p)$.

In order to fix an interpretation of linear logic in a net we just need an interpretation of the atomic propositions as elements of the associated quantale. For simplicity we consider a linear logic language where the atomic propositions a name singleton multisets of places \underline{a} , not necessarily injectively. The net, together with the naming function, we call a *net valuation*. Formulae are given by:

$$\begin{array}{ll}
A ::= \mathbf{T} \mid \mathbf{F} \mid \mathbf{1} & \text{constants} \\
\mid a & \text{atomic propositions} \\
\mid A \otimes A \mid A \multimap A & \text{multiplicative connectives} \\
\mid A \& A \mid A \oplus A & \text{additive connectives}
\end{array}$$

We make the choice of interpreting an atomic proposition as the downwards closure of the associated place, but we could just as well have used the downwards closure of some marking without altering our results. This choice is consistent with the following intuitive understanding: the denotation of an assertion is to be thought of as the set of requirements sufficient to establish it.

With respect to a net valuation N , linear logic formulae are interpreted as follows:

$$\begin{array}{ll}
\llbracket \mathbf{T} \rrbracket_N & = \mathcal{M} \\
\llbracket \mathbf{F} \rrbracket_N & = \emptyset \\
\llbracket \mathbf{1} \rrbracket_N & = \{M \mid M \rightarrow \underline{\mathbf{0}}\} \\
\llbracket a \rrbracket_N & = \{M \mid M \rightarrow \underline{a}\} \\
\llbracket A \otimes B \rrbracket_N & = \{M \mid \exists M_A \in \llbracket A \rrbracket_N, M_B \in \llbracket B \rrbracket_N. M \rightarrow M_A + M_B\} \\
\llbracket A \multimap B \rrbracket_N & = \{M \mid \forall M_A \in \llbracket A \rrbracket_N. M + M_A \in \llbracket B \rrbracket_N\} \\
\llbracket A \& B \rrbracket_N & = \llbracket A \rrbracket_N \cap \llbracket B \rrbracket_N \\
\llbracket A \oplus B \rrbracket_N & = \llbracket A \rrbracket_N \cup \llbracket B \rrbracket_N
\end{array}$$

The semantics of [Bro89] is similar, but somehow dual to that here.

Because of the interpretation of $\mathbf{1}$ validity of an assertion A for the given net valuation, N , can be expressed by:

$$\models_N A \quad \text{iff} \quad \underline{\mathbf{0}} \in \llbracket A \rrbracket_N .$$

Semantic entailment between assertions A and B amounts to:

$$A \models_N B \quad \text{iff} \quad \llbracket A \rrbracket_N \subseteq \llbracket B \rrbracket_N .$$

Because of the interpretation of linear implication, this is equivalent to

$$\models_N A \multimap B .$$

General validity, $\models A$, of an assertion A is defined by

$$\models A \quad \text{iff} \quad \models_N A, \text{ for every net valuation } N$$

and with respect to entailment: $\Gamma \models B$ iff $\Gamma \models_N B$, for every net valuation N .

As a special case that quantale semantics is sound, we have the soundness result:

Theorem 6 *If $\Gamma \vdash A$ then $\Gamma \models A$.*


So we see that with respect to a Petri net, an assertion A is denoted a set of markings $\llbracket A \rrbracket_N$. As we have discussed, a marking of net can be viewed as a distribution of resources. When $M \in \llbracket A \rrbracket_N$ we can think of the marking M as a distribution of resources sufficient to establish A according to the net; in this sense the marking M is one of the (in general many) requirements sufficient to establish A . The meaning of an assertion A is specified by saying what requirements are sufficient to establish it—this is the content of the denotation $\llbracket A \rrbracket_N$. Accordingly, a net satisfies an assertion A when $\underline{0} \in \llbracket A \rrbracket_N$, expressing that A can be established with no resources.

This reading squares with the fact that assertions denote subsets of markings which are downwards closed with respect to the reachability relation of the net; if $M \in \llbracket A \rrbracket_N$, so M is a requirement sufficient to establish A , and $M' \rightarrow M$ so we can obtain M for M' , then so also is M' a sufficient requirement of A . Casting an eye over the definition of the semantics of assertions we can read, for example, the definition of $\llbracket a \rrbracket_N$, for an atom a , as expressing that a sufficient requirement of a is any marking from which the marking \underline{a} can be reached according to the net. Similarly, the sufficient requirements of $A \& B$ are precisely those which are sufficient requirements of both A and of B . An element of $\llbracket A \multimap B \rrbracket_N$ can be seen as what is required, in addition to any requirement of A , in order to establish B . There are similar restatements of the semantics for the other connectives as well.

This understanding should be born in mind when considering the examples that follow, where we shall make use of the fact that \otimes , $\&$ and \oplus are associative and assume the precedence:

$$\multimap < \&, \oplus < \otimes .$$


Consider the net:



Here we have $\llbracket b \rrbracket_{N_1} = \{\underline{d}, \underline{a}, \underline{b}\}, \dots, \llbracket c \rrbracket_{N_1} = \{\underline{a}, \underline{c}\}$ and $\llbracket b \otimes c \rrbracket_{N_1} = \{\underline{d} + \underline{a}, \underline{d} + \underline{c}, \underline{a} + \underline{a}, \underline{a} + \underline{c}, \underline{b} + \underline{a}, \underline{b} + \underline{c}\}$, so consequently

$$\begin{aligned} \llbracket d \otimes a \rrbracket_{N_1} &\subseteq \llbracket b \otimes c \rrbracket_{N_1} \text{ or equivalently } \models_{N_1} d \otimes a \multimap b \otimes c \\ \models_{N_1} a \otimes a &\multimap b \otimes c \\ \models_{N_1} a &\multimap b \oplus c, \models_{N_1} d \oplus a \multimap b, \models_{N_1} d \multimap b \oplus c \\ \models_{N_1} b \&c \multimap e, \models_{N_1} a &\multimap b \&c \quad . \end{aligned}$$


The most difficult connective to comprehend is probably linear implication so we give a few more examples here. For the nets




we have:

$$\begin{aligned} \llbracket c \multimap d \rrbracket_{N_2} &= \emptyset \text{ and } \models_{N_2} (c \multimap d) \multimap \mathbf{F} \\ \llbracket b \multimap d \rrbracket_{N_2} &= \{\underline{a}\} \text{ and } \models_{N_2} (b \multimap d) \multimap a \\ \llbracket a \multimap e \rrbracket_{N_3} &= \{\underline{b} + \underline{d}\} \quad . \end{aligned}$$

A more peculiar example of linear implication is given by the following net N :



Exercise 7 Determine $\llbracket d \rrbracket_N$, $\llbracket d \otimes e \rrbracket_N$, $\llbracket d \& e \rrbracket_N$, $\llbracket b \multimap d \rrbracket_N$ and $\llbracket c \multimap d \rrbracket_N$ for:



□

Exercise 8

(i) Construct a net N which shows that the distributivity

$$(A \otimes C) \& (B \otimes C) \models_N (A \& B) \otimes C$$

fails.

(ii) Prove that $(A \& B) \otimes C \models (A \otimes C) \& (B \otimes C)$.

(iii) Does $(A \& C) \otimes (B \& C) \models (A \otimes B) \& C$ hold?

(iv) What about $(A \otimes B) \& C \models (A \& C) \otimes (B \& C)$? □

Exercise 9 Which of the following statements are true, if any?

(i) $A \oplus (B \otimes C) \models (A \oplus B) \otimes (A \oplus C)$

(ii) $(A \oplus B) \otimes (A \oplus C) \models A \oplus (B \otimes C)$ □

8 Expressing Properties of Nets

Notation: For a multiset, M , of assertions of our logic, we associate the formula \widehat{M} which when M is nonempty is given by

$$\bigotimes_{M(A) \neq 0} A^{M(A)} \quad \text{where } A^n = \overbrace{A \otimes \cdots \otimes A}^n, \text{ for } n > 0$$

and otherwise, when $M = \underline{0}$, is given by the formula **1**. With respect to a net valuation, for a multiset M of atoms of the logic we can write \underline{M} for the marking satisfying

$$\underline{M}(p) = \sum_{a=p} M(a)$$

We let the context distinguish M , and its understanding as \widehat{M} and \underline{M} , except for a few crucial statements and proofs.

We can then express that one marking is reachable in a net N :

Proposition 10 *For any multisets of atoms M and M' ,*

$$M \rightarrow M' \text{ in the net } N \quad \text{iff} \quad \models_N M \multimap M' .$$

Proof Simply note that $M \rightarrow M'$ in N iff $\downarrow M \subseteq \downarrow M'$ iff $\llbracket M \rrbracket_N \subseteq \llbracket M' \rrbracket_N$ iff $\models_N M \multimap M'$. \square

Before reading the list of sample properties below observe:

$$\llbracket M \otimes \mathbb{T} \rrbracket_N = \downarrow \{M' \mid M' \geq M\} .$$


Sample Properties

Suppose a net N with initial marking M_0 .

- From the initial marking it is possible to reach a marking where a is marked:
 $\models_N M_0 \multimap a \otimes \mathbb{T}$.
- From the initial marking it is possible to reach a marking where a place, a , of $S \subseteq^{fn} P$ is marked: $\models_N M_0 \multimap (\bigoplus_{a \in S} a) \otimes \mathbb{T}$.
- Once a is marked it is possible to reach a marking where b is marked:
 $\models_N a \otimes \mathbb{T} \multimap b \otimes \mathbb{T}$.
- Once a is marked either b or c can become marked (but we do not know which): $\models_N a \otimes \mathbb{T} \multimap (b \oplus c) \otimes \mathbb{T}$.
- Once a is marked both b and c can become marked (but not necessarily simultaneously): $\models_N a \otimes \mathbb{T} \multimap (b \& c) \otimes \mathbb{T}$.

Example 11 (Mutual exclusion)

Consider the net N :



outside its critical region, c_1 , and similarly for the other process, p_2 . The resource corresponding to b is used to ensure mutual exclusion of the critical regions and after a process has been in its critical region it returns a resource, a , which then is prepared (transformed into b) for the next turn. The initial marking, M_0 , will be $M_0 = b_0 \otimes w_1 \otimes w_2$. We can now express that e.g. p_1 can enter its critical region (from the initial marking) by: $\models_N M_0 \multimap c_1 \otimes \mathbf{T}$. However this does not ensure that no undesired tokens are present, so it is better to express it: $\models_N M_0 \multimap c_1 \otimes w_2$. If the system is in a “working state” then both processes have the possibility of entering their critical section: $\models_N w_1 \otimes (a \oplus b) \otimes w_2 \multimap c_1 \otimes w_2 \ \& \ w_1 \otimes c_2$. The property, that when p_1 is in its critical section and p_2 is working it is possible that p_2 can later come into its critical section with p_1 working, is expressed by: $\models_N c_1 \otimes w_2 \multimap w_1 \otimes c_2$. Similar other “positive” properties can be expressed. Shortly we shall see how to express the “negative” property that both processes cannot be in their critical regions at the same time.

Exercise 12 Let N be the net of example 11.

(i) Which of the following statements are true?


$$\models_N (c_1 \multimap w_1) \multimap \mathbf{F}, \quad \models_N (w_1 \multimap c_1) \multimap \mathbf{F}, \quad \models_N (M_0 \multimap c_1 \otimes c_2) \multimap \mathbf{F}$$

(ii) A new proof system for the net N can be obtained by adding to the proof rules an axiom $\widehat{\bullet}t \vdash \widehat{\bullet}t$ for each transition t in the net, e.g.

$$\frac{}{w_1 \otimes b \vdash c_1}.$$

Prove $\vdash w_1 \otimes (a \oplus b) \otimes w_2 \multimap c_1 \otimes w_2$ in this new proof system. □

Exercise 13 Consider the following net N_V describing a vending machine for cheap hot drinks:



Express in linear logic the following statements and consider in what way the statements are made more precise.

- (ii) For 2 pence one can get a cup of coffee and a cup of tea.
- (iii) For five pence one can get a cup of coffee and in addition a cup of tea as well as a penny back.
- (iv) If one has a penny and a part from that sufficient money to get both tea and coffee then one can also get chocolate.
- (v) Think of more yourself. □

9 The “Of Course” Operator

From the discussion in section 4 it immediately follows how to interpret an “of course” formula:

Definition 14 *Given a net N with associated quantale \mathcal{Q} and a formula A , the interpretation of “of course” A , is*

$$\llbracket !A \rrbracket_N = \bigcup \{q \in \mathcal{Q} \mid q \text{ is a postfix point of } f_A\}$$

where $f_A : \mathcal{Q} \rightarrow \mathcal{Q}$ is the function given by:

$$x \mapsto \llbracket \mathbf{1} \rrbracket_N \cap \llbracket A \rrbracket_N \cap (x \otimes x) \tag{9}$$

□

In order to gain a better understanding of the “of course” operator we give the following characterisation of a postfix point.

Proposition 15 *$q \in \mathcal{Q}$ is a post fixed point of (9) iff there exists an $H \subseteq \mathcal{M}$ such that*

- (i) $q = \downarrow H$
- (ii) $H \subseteq \llbracket A \rrbracket_N$
- (iii) $\forall M \in H. M \rightarrow \underline{\mathcal{Q}}$
- (iv) $\forall M \in H \exists M', M'' \in H. M \rightarrow M' + M''$.


Proof if: $\downarrow H = q$ a postfix point of (9) means $q \subseteq \mathbf{1} \cap \llbracket A \rrbracket_N \cap (q \otimes q)$. From (iii) follows $q = \downarrow H \subseteq \downarrow \underline{\mathcal{Q}} = \mathbf{1}$ and from (ii) $q = \downarrow H \subseteq \downarrow \llbracket A \rrbracket_N = \llbracket A \rrbracket_N$ as $\llbracket A \rrbracket_N$ is downwards closed. To see $q \subseteq q \otimes q$ assume $M \in q$. Then $\exists M''' \in H. M \rightarrow M'''$

$M \in \downarrow(H + H) = H \otimes H \subseteq \downarrow H \otimes \downarrow H = q \otimes q$.

only if: Take $H = q$ and we have $\downarrow H = H$. Properties (i) – (iii) follows readily as does (iv) by $q \subseteq q \otimes q$ and the downwards closure of H . \square

In the light of this proposition, we can see a requirement of $!A$ as any requirement of A which can both vanish and duplicate into any number of additional requirements of $!A$.

Example 16 For the net N :



we get: $\llbracket b \rrbracket_N = \{\underline{b}\} \cup \{n\underline{a} \mid n > 0\}$ which properly contains $\llbracket !b \rrbracket_N = \{n\underline{a} \mid n > 0\}$.

10 Linear Negation

We have discovered that in a net N one useful interpretation of the extra logical constant \perp , for linear-absurdity, is:

$$\llbracket \perp \rrbracket_N = \{M \mid \underline{0} \not\rightarrow M \text{ in } N\} .$$

That is \perp denotes exactly the set of markings which cannot be reached from the empty marking. For $\llbracket \perp \rrbracket_N$ to be well-defined, we need that it is downwards-closed: To see this, suppose $M' \rightarrow M$ for $M \in \llbracket \perp \rrbracket_N$. Assume $M' \notin \llbracket \perp \rrbracket_N$. Then $\underline{0} \rightarrow M'$, whence from $M' \rightarrow M$ and transitivity of \rightarrow we conclude $\underline{0} \rightarrow M$ —contradicting $M \in \llbracket \perp \rrbracket_N$.

Since linear negation can be expressed in terms of \perp and \multimap (by $A^\perp = A \multimap \perp$), we get:

$$\llbracket A^\perp \rrbracket_N = \{M \mid \forall M_A \in \llbracket A \rrbracket_N. \underline{0} \not\rightarrow M_A + M\} .$$

The interesting consequence of this particular choice for \perp is, that whatever property we could state before in terms of validity of a formula A can now be stated negatively as $\models_N A^\perp$. Formally:

Proposition 17

$$\models_N A^\perp \quad \text{iff} \quad \not\models_N A \tag{10}$$

$\underline{Q} \not\rightarrow M$ in N iff $\exists M_A \in \llbracket A \rrbracket_N. \underline{Q} \rightarrow M_A$ iff $\underline{Q} \in \llbracket A \rrbracket_N$ iff $\models_N A$, where the second last equivalence follows from the downwards closure of $\llbracket A \rrbracket_N$. \square

Clearly $\{M \mid \underline{Q} \not\rightarrow M \text{ in } N\}$ is the largest element which can be used for the interpretation of \perp should (10) hold. Suppose on the contrary $M \in \llbracket \perp \rrbracket_N$ and $M \notin \{M \mid \underline{Q} \not\rightarrow M \text{ in } N\}$. Then $\underline{Q} \rightarrow M$ and so $\models_N \perp$ but also $\models_N \perp^\perp$.

If furthermore each place is named by some atomic proposition then (10) uniquely determines the interpretation of \perp . To see this notice that each marking is then denotable and that

$$\begin{aligned} \underline{M} \in \llbracket \perp \rrbracket_N &\text{ iff } \widehat{M} \models_N \perp \\ &\text{ iff } \models_N \widehat{M} \multimap \perp \\ &\text{ iff } \models_N \widehat{M}^\perp \\ &\text{ iff } \not\models_N \widehat{M} \text{ by (10)} \\ &\text{ iff } \underline{Q} \not\rightarrow \underline{M} \text{ in } N \end{aligned}$$

Combining proposition 17 with proposition 10 we can express that a marking M' cannot be reached from another M :

Corollary 18 *For markings M and M' :*

$$M \not\rightarrow M' \quad \text{iff} \quad \models_N (M \multimap M')^\perp .$$

Example 11 (continued)

We can now express that the processes, p_1 and p_2 cannot get into their critical regions at the same time. We might try $\models_N (M_0 \multimap c_1 \otimes c_2)^\perp$. This is not quite right however, since $\models_N (M_0 \multimap c_1 \otimes c_2)^\perp$ merely states that the two processes cannot be in their critical regions at the same time when no other tokens are present; the correct statement is $\models_N (M_0 \multimap c_1 \otimes c_2 \otimes \mathbf{T})^\perp$.

Sample Properties

Suppose a net N .

- There is a marking with a marked such that b will never be marked in any reachable marking:

$$\models_N (a \otimes \mathbf{T} \multimap b \otimes \mathbf{T})^\perp .$$

- For any marking obtained from M the *load* on the place a cannot exceed $n \in \mathbb{N}$, i.e. with $M = M_0$ this means that a is *n-safe*:

$$\models_N (M \multimap a^{n+1} \otimes \mathbf{T})^\perp .$$

Then M' belongs to $\llbracket a_N^{n+1} \otimes \mathbf{T} \rrbracket$ which is downwards closed and so $M \in \llbracket a^{n+1} \otimes \mathbf{T} \rrbracket_N$ wherefore $\models_N M \multimap a^{n+1} \otimes \mathbf{T}$ —contradicting $\models_N (M \multimap a^{n+1} \otimes \mathbf{T})^\perp$.

- That a transition t is *M-dead* in a net N , i.e. $\forall M' \in [M]. M' \not\{t\}$, is expressed by:

$$\models_N (M \multimap \bullet t \otimes \mathbf{T})^\perp .$$

- If N only have finitely many places, we can express *safeness*, i.e. $\forall M \in [M_0] \forall a \in P. M(a) \leq 1$ (N is *1-safe*):

$$\models_N (M_0 \multimap (\bigoplus_{a \in P} a \otimes a) \otimes \mathbf{T})^\perp .$$

Again suppose on the contrary $\exists M' \exists a. M_0 \rightarrow M'$ and $M'(a) > 1$. Then $M' \in \llbracket (\bigoplus_{a \in P} a \otimes a) \otimes \mathbf{T} \rrbracket_N$ and so $\models_N M_0 \multimap (\bigoplus_{a \in P} a \otimes a) \otimes \mathbf{T}$ —a contradiction.

This definition of linear negation actually allows us to state any propositional combination of validity of linear formulae simply as the validity of a single formula. Abbreviating in a formula $A^\perp \oplus B$ by $A \Rightarrow B$ we can read \Rightarrow as implication, \oplus as disjunction etc. For example:

if b can be reached from a then so can c :

$$\models_N a \multimap b \text{ implies } \models_N a \multimap c$$

can be expressed by a single formula:

$$\models_N (a \multimap b) \Rightarrow (a \multimap c)$$

Exercise 19 Formulate the following statement in linear logic.

If it is possible to reach b from a in the net then c can be reached as well. \square

It follows from the remarks of section 4 that by denoting a proposition A by $\llbracket A \rrbracket_N^{\perp\perp}$ we obtain a Petri-net semantics of classical linear logic. In general it is hard to get an impression of $\llbracket A \rrbracket_N^{\perp\perp}$, but by slightly narrowing the class of nets we consider to those whose transitions always have a nonempty preset, i.e. $\underline{0} \neq \bullet t$ for all transitions t , we get:

$$\llbracket A \rrbracket_N^{\perp\perp} = \begin{cases} \llbracket \mathbf{T} \rrbracket_N & \text{if } \models_N A \\ \llbracket \mathbf{F} \rrbracket_N & \text{otherwise} \end{cases} . \quad (11)$$

To see this notice first for each net we have $\llbracket \perp \rrbracket_N = \{M \mid M \neq \underline{0}\} = \mathcal{M} \setminus \{\underline{0}\}$ and so

$$\begin{aligned} q^\perp &= \{M \mid \forall M_q \in q. M + M_q \neq \underline{0}\} \\ &= \mathcal{M} \setminus \{M \mid \exists M_q \in q. M + M_q = \underline{0}\} \\ &= \begin{cases} \mathcal{M} \setminus \{\underline{0}\} & \text{if } \underline{0} \in q \\ \mathcal{M} & \text{otherwise} \end{cases} . \end{aligned}$$

a 2-element quantale where the logical operations act as the usual operations of propositional logic, and \otimes acts like $\&$. In this quantale all the atomic propositions would have the same interpretation as \mathbf{F} and \perp . Such a classical interpretation seems of limited interest.

11 Recursion

Though we now have the possibility to use linear negation to express properties of a net it is hard to express properties such as:

From the initial marking it is possible to reach a marking where just a is marked (to arbitrary multiplicity).

One way to increase the power of the logic language is to extend it with the possibility of writing recursive formulae using least (μ) and greatest (ν) fixpoint operators on assertions.

With their aid we can define propositions a^+ and a^* . Abbreviating $\mu x. a \oplus (a \otimes x)$ by a^+ and $\mathbf{1} \oplus a^+$ by a^* we have:

$$\begin{aligned} M \in \llbracket a^+ \rrbracket & \quad \text{iff} \quad \exists n > 0. M \rightarrow n\underline{a} \\ M \in \llbracket a^* \rrbracket & \quad \text{iff} \quad \exists n \geq 0. M \rightarrow n\underline{a} \end{aligned}$$

Clearly the “of course” operator can be expressed as a maximum fixed point: $!A = \nu x. A \& \mathbf{1} \& x \otimes x$.

Sample Properties

Suppose a net N .

- From the initial marking it is possible to reach a marking where just a is marked: $\models_N M_0 \multimap a^+$.
- From the marking M it is possible in a finite net to reach a marking with a marked and b unmarked:
 $\models_N M \multimap a^+ \otimes (\otimes_{c \in P \setminus \{a,b\}} c^*)$.
- The places of $S \subseteq^{fn} P$ can always be emptied of tokens: $\models_N \mathbf{T} \multimap \otimes_{a \in P \setminus S} a^*$.

Another possible extension is to add second order quantifiers with variables ranging over elements of the quantale, or quantifiers with variables over markings, for example; they can be understood as special kinds of meet and join operations. We introduce a restricted form of second-order quantification in section 14.


Linear logic induces an equivalence relation between nets having the same set of places. With respect to a set of places, we build up the propositions of a linear logic with the places as atoms. If A is such a proposition whose atoms are amongst the places of a net N we say N satisfies A , written $\models_N A$, iff A is valid with respect to the semantics of section 7. We say two nets N_0, N_1 with the same places are equivalent if they make the same propositions valid, i.e. we define

$$N_0 \sim N_1 \quad \text{iff} \quad (\models_{N_0} A \text{ iff } \models_{N_1} A, \text{ for all propositions } A) .$$

A less refined equivalence can be got by taking two nets to be equivalent iff they give rise to the isomorphic quantales. The latter notion has the advantage of not depending on the nets having the same set of places. We shall concentrate on the former equivalence as it implies the latter and suffices to point out some of the idiosyncrasies of linear logic when interpreted on nets. The equivalence has a direct bearing on the suitability of linear logic as a specification language for Petri nets. If it is wished to distinguish between two nets which are equivalent, there is no hope of specifying one and not the other within pure linear logic.

Observe that because our semantics of linear logic is built on the relation \rightarrow , any two nets, over a common set of places, which induce the same relation \rightarrow are bound to be equivalent. This means that whenever $M \rightarrow M'$, for a net N , we can adjoin a new transition t with the pre places M and post places M' , and still get a net equivalent to N . So, given any net N , we can add “identity” transitions at any multiset of places, “compositions” and “synchronizations” of transitions as new transitions and obtain a net equivalent to N .

Examples



Also note that these two nets are equivalent:



where the direction of the loop has changed! (It’s easy to see both nets have the

looping if another place is present to act like a loop counter:




These nets are not equivalent.

Notice that whether or not the equivalence induced on nets is reasonable, if two nets are equivalent in our sense then so are they too with respect to the satisfaction relation of [MOM89]. This is because their construction of a linear category from a net factors through the construction of what they call a Petri-category. The Petri-category made from a net has objects the net's markings and has a morphism $f : M \rightarrow M'$ iff $M \rightarrow M'$ in our sense. A difference with the approach of [MOM89] can be seen by considering a preorder induced by the logic rather than just equivalence. For nets with the same set of places, define

$$N_0 \preceq N_1 \quad \text{iff} \quad (\models_{N_0} A \text{ implies } \models_{N_1} A, \text{ for all propositions } A) .$$

The preorder induced by our definition of satisfaction will differ from theirs. In [MOM89], if a net N satisfies a formula A , in their sense, then any net obtained from N by adding an arbitrary number of new transitions will also satisfy A . This is not the case for our definition of satisfaction, as can be seen by adding transitions to the nets N_1 and N_2 in section 7: Take



For these nets we have $\not\models_{N'_1} b \& c \multimap e$ and $\not\models_{N'_2} (b \multimap d) \multimap a$ whereas $\models_{N_1} b \& c \multimap e$ and $\models_{N_2} (b \multimap d) \multimap a$.

These remarks imply that proof system described in (ii) of exercise 12 does not describe a net up to \sim .

In this section we shall be concerned with completeness results for different fragments of linear logic without exponentials.

We start by sketching the completeness proof for quantale semantics.

The idea in showing completeness is to build a quantale by taking the ideal completion of the Lindenbaum algebra. More precisely take Q to be \subseteq -ordered set of subsets I of assertions of linear logic, without exponentials, such that

$$\begin{aligned} A \vdash B \in I &\Rightarrow A \in I, \\ X \subseteq^{fn} I &\Rightarrow \oplus X \in I \\ (\text{We understand } \oplus \emptyset &= \mathbf{F}) \quad . \end{aligned}$$

The \otimes -operation on ideals is got by taking:

$$I \otimes J =_{def} \{C \mid \exists A \in I, B \in J. C \vdash A \otimes B\} .$$

That this yields an ideal follows routinely: clearly $I \otimes J$ is closed with respect to entailment, *i.e.* $B \vdash C \in I \otimes J$ implies $B \in I \otimes J$; it is closed under \oplus because it contains $\oplus \emptyset = \mathbf{F}$ and if $C, C' \in I \otimes J$ then $C \vdash A \otimes B$ and $C' \vdash A' \otimes B'$, for $A, A' \in I, B, B' \in J$, whence

$$\begin{aligned} C \oplus C' &\vdash (A \otimes B) \oplus (A' \otimes B') \\ &\vdash (A \otimes B) \oplus (A' \otimes B) \oplus (A \otimes B') \oplus (A' \otimes B') \\ &\vdash (A \oplus A') \otimes (B \oplus B'), \end{aligned}$$

where $(A \oplus A') \in I$ and $(B \oplus B') \in J$ thus ensuring $C \oplus C' \in I \otimes J$ —thus it is closed under \oplus .

The quantale Q interprets assertions once we decide to interpret atoms a in the following way:

$$\llbracket a \rrbracket_Q = \{B \mid B \vdash a\} .$$

It is a relatively simple matter to show the following agreement between the semantics in the constructed quantale and the proof system:

Lemma 20 *Letting A be an assertion of linear logic without exponentials,*

$$\llbracket A \rrbracket_Q = \{B \mid B \vdash A\} .$$

Proof By structural induction. We consider two cases:

$A \equiv A_1 \oplus A_2$: The denotation $\llbracket A_1 \oplus A_2 \rrbracket_Q = \llbracket A_1 \rrbracket_Q \vee \llbracket A_2 \rrbracket_Q$, the join in Q , which contains $A_1 \oplus A_2$, and hence must equal the principal ideal $\{B \mid B \vdash A_1 \oplus A_2\}$.

$$\llbracket A_1 \multimap A_2 \rrbracket_Q = \bigvee \{ I \mid I \otimes \llbracket A_1 \rrbracket_Q \subseteq \llbracket A_2 \rrbracket_Q \}$$

a join in Q which contains $A_1 \multimap A_2$ and hence includes the principal ideal $\{B \mid B \vdash A_1 \multimap A_2\}$. It is in fact equal to this principal ideal. To see this, let $B \in I$ where $I \otimes \llbracket A_1 \rrbracket_Q \subseteq \llbracket A_2 \rrbracket_Q$. Then $B \otimes A_1 \in \llbracket A_2 \rrbracket_Q$, so by structural induction $B \otimes A_1 \vdash A_2$, whence $B \vdash A_1 \multimap A_2$. \square

Corollary 21 *Let A be an assertion of linear logic without exponentials. Then*

$$\models_Q A \quad \text{iff} \quad \vdash A .$$

As a corollary we from $A \vdash B$ iff $\vdash A \multimap B$ obtain completeness with respect to quantales:

Theorem 22 *For the fragment without exponentials we have:*

$$\Gamma \models A \quad \text{iff} \quad \Gamma \vdash A .$$

In the remaining sections we shall only be concerned with completeness proofs for the net semantics.

Completeness for the \oplus -free Fragment


Restrict the syntax to the fragment:

$$A ::= \top \mid \mathbf{1} \mid a \mid A_1 \& A_2 \mid A_1 \otimes A_2 \mid A_1 \multimap A_2 \quad (\oplus\text{-free})$$

where a ranges over atoms. For the \oplus -free fragment we construct a net N where the places are formulae and the transitions essentially correspond to the provable sequents. I.e.

- *Places* are assertions of $(\oplus\text{-free})$ above. An atom will be interpreted as its singleton marking.
- *Transitions* are pairs (M, M') of multisets of places for which $\widehat{M} \vdash \widehat{M}'$ with pre- and postset maps $\bullet(M, M') = M$ and $(M, M')\bullet = M'$.

For instance the net has the following transition



Lemma 23 For markings M, M' of the net N ,

$$M \rightarrow M' \text{ in the net} \quad \text{iff} \quad \widehat{M} \vdash \widehat{M}' .$$

Proof “if”: It is clear by definition that if $\widehat{M} \vdash \widehat{M}'$ then $M \rightarrow M'$ for any markings $M, M' \in \mathcal{M}$.

“only if”: Follows by a simple inductive argument once we have established

$$M [t] M' \text{ implies } \widehat{M} \vdash \widehat{M}' .$$

However, if $M [t] M'$ then, by definition, there is some $M'' \in \mathcal{M}$ such that

$$M = M'' + \bullet t \text{ and } t \bullet + M'' = M' .$$

From $\widehat{\bullet t} \vdash \widehat{t \bullet}$ we derive $\widehat{M}'' \otimes \widehat{\bullet t} \vdash \widehat{t \bullet} \otimes \widehat{M}''$. The result then follows from $\widehat{M} \dashv\vdash \widehat{M}'' \otimes \widehat{\bullet t}$ and $\widehat{M}' \dashv\vdash \widehat{t \bullet} \otimes \widehat{M}''$. \square

Lemma 24 For the \oplus -free fragment we have: $\llbracket A \rrbracket_N = \{M \mid \widehat{M} \vdash A\}$.

Proof By induction on the structure of A using the previous lemma.

$A \equiv \top$: $\llbracket \top \rrbracket_N = \mathcal{M} = \{M \in \mathcal{M} \mid M \vdash \top\}$ by axiom $\Gamma \vdash \top$ (recall \mathcal{M} consists of finite multisets).

$A \equiv \mathbf{1}$: $\llbracket \mathbf{1} \rrbracket_N = \{M \in \mathcal{M} \mid M \rightarrow \mathbf{0}\} = \{M \in \mathcal{M} \mid M \vdash \widehat{\mathbf{0}} = \mathbf{1}\}$ by lemma 23.

$A \equiv a$: $\llbracket a \rrbracket_N = \{M \in \mathcal{M} \mid M \rightarrow a\} = \{M \in \mathcal{M} \mid M \vdash a\}$ by lemma 23.

$A \equiv A_1 \otimes A_2$: $M \in \llbracket A_1 \otimes A_2 \rrbracket_N$
 $\Leftrightarrow \exists M_1 \in \llbracket A_1 \rrbracket_N, M_2 \in \llbracket A_2 \rrbracket_N. M \rightarrow M_1 + M_2$ by definition,
 $\Leftrightarrow \exists M_1, M_2 \in \mathcal{M}. M_1 \vdash A_1, M_2 \vdash A_2$ and $M \rightarrow M_1 + M_2$ by induction,
 $\Leftrightarrow \exists M_1, M_2 \in \mathcal{M}. M_1 \vdash A_1, M_2 \vdash A_2$ and $M \vdash M_1 \otimes M_2$ by lemma 23,
 $\Leftrightarrow M \vdash A_1 \otimes A_2$ by $(\vdash \otimes)$, $(\otimes \vdash)$ and (identity).

$A \equiv A_1 \multimap A_2$:

$M \in \llbracket A_1 \multimap A_2 \rrbracket_N \Leftrightarrow \forall M_1 \in \llbracket A_1 \rrbracket_N. M + M_1 \in \llbracket A_2 \rrbracket_N$ by definition,
 $\Leftrightarrow \forall M_1 \in \mathcal{M}. M_1 \vdash A_1 \Rightarrow M \otimes M_1 \vdash A_2$ by induction,
 $\Rightarrow M \vdash A_1 \multimap A_2$ by $(\vdash \multimap)$ and (identity).

To see “ \Leftarrow ” suppose $M \vdash A_1 \multimap A_2$. From $(\multimap \vdash)$ we derive $M, A_1 \vdash A_2$. Using (cut) and $M_1 \vdash A_1$ we then get $M, M_1 \vdash A_2$.

$A \equiv A_1 \& A_2$: $M \in \llbracket A_1 \& A_2 \rrbracket_N \Leftrightarrow M \in \llbracket A_1 \rrbracket_N$ and $M \in \llbracket A_2 \rrbracket_N$ by definition,
 $\Leftrightarrow M \vdash A_1$ and $M \vdash A_2$ by induction,
 $\Rightarrow M \vdash A_1 \& A_2$ by $(\vdash \&)$.

For the other direction “ \Leftarrow ” we by (identity) and $(l\&\vdash)$ obtain $A_1 \& A_2 \vdash A_1$ and so $M \vdash A_1$ from $M \vdash A_1 \& A_2$ and (cut). By symmetry $M \vdash A_2$. \square

duce:

Theorem 25 *For the \oplus -free fragment we have:*

$$\Gamma \models A \quad \text{iff} \quad \Gamma \vdash A .$$

As observed by Sergei Soloviev, the net need for a particular sequent only to be constructed with a finite number of places corresponding to subformulae of the sequent. However, it not clear that the net can be finite if the sequent contains $\&$ or \multimap .

Completeness for the \multimap -free Fragment

We can obtain completeness for the \multimap -free fragment of propositional intuitionistic logic. Its syntax:

$$A ::= \mathbf{T} \mid \mathbf{F} \mid \mathbf{1} \mid a \mid A_1 \oplus A_2 \mid A_1 \& A_2 \mid A_1 \otimes A_2 \quad (\multimap\text{-free})$$

where a ranges over atoms. With a similar construction to that in the previous section we can define a net N (this time with places in the \multimap -free fragment) to obtain a rather weak form of completeness for the \multimap -free fragment.

Lemma 26 *For the \multimap -free fragment we have $\llbracket A \rrbracket_N \subseteq \{M \mid M \vdash A\}$.*

Proof Induction on the structure of A . All the cases except $A \equiv \mathbf{F}$ and $A \equiv A_1 \oplus A_2$ are handled exactly as the \subseteq -part of lemma 24 (notice the weaker hypothesis).

$A \equiv \mathbf{F}$: Evident as $\llbracket \mathbf{F} \rrbracket_N = \emptyset$.

$A \equiv A_1 \oplus A_2$: $M \in \llbracket A_1 \oplus A_2 \rrbracket_N \Leftrightarrow M \in \llbracket A_1 \rrbracket_N$ or $M \in \llbracket A_2 \rrbracket_N$ by definition,
 $\Rightarrow M \vdash A_1$ or $M \vdash A_2$ by induction,
 $\Rightarrow M \vdash A_1 \oplus A_2$ by $(\vdash \oplus l)$ or $(\vdash \oplus r)$. □

As a corollary we have:

Theorem 27 *For the \multimap -free fragment we have:*

$$\models A \quad \text{iff} \quad \vdash A .$$

We have not used the distributive law yielded by the net semantics:

$(A \oplus B) \& C \vdash (A \& C) \oplus (B \& C) \quad (\&\text{-}\oplus\text{-dist.})$

for the \multimap -free fragment of propositional intuitionistic logic.

To show completeness we construct a net with places (and markings) identified with assertions in the \oplus -free subfragment:

$$A ::= \top \mid \mathbf{1} \mid a \mid A_1 \& A_2 \mid A_1 \otimes A_2 \quad (\multimap\text{-}\oplus\text{-free})$$

We will just call it the \oplus -free fragment in the rest of this section. Construct a net N where:

- *Places* are assertions in the \oplus -free fragment (atoms are to be interpreted as their singleton markings).
- *Transitions* are pairs (M, M') of multisets of places for which $\widehat{M} \vdash \widehat{M}'$.

Lemma 28 *For markings M, M' of the net,*

$$M \rightarrow M' \text{ in the net} \quad \text{iff} \quad M, M' \text{ } \oplus\text{-free and } M \vdash M' \text{ in the logic.}$$

Proof The proof is like that for lemma 23. □

Lemma 29 (Decomposition lemma). *For any \multimap -free assertion A there is a finite set I indexing $\multimap\text{-}\oplus\text{-free}$ assertions M_i , such that*

$$A \dashv\vdash \bigoplus_{i \in I} M_i \text{ .}$$

Proof The proof proceeds by structural induction on the assertion A .

The base cases are routine; for example $\mathbf{F} \dashv\vdash \bigoplus \emptyset$ ($= \mathbf{F}$ by definition), i.e. falsity is interderivable with the empty disjunction. Of the remaining cases, that where A has the form $A_1 \& A_2$ makes use, as is to be expected, of the additional distributivity rules for $\&$ and \oplus . Inductively, assume

$$A_1 \dashv\vdash \bigoplus_{i \in I} M_i^1 \quad \text{and} \quad A_2 \dashv\vdash \bigoplus_{j \in J} M_j^2 \text{ .}$$

Then, from these assumptions and repeated use of $\&\text{-}\oplus\text{-distributivity}$

$$\begin{aligned} A_1 \& A_2 &\dashv\vdash (\bigoplus_i M_i^1) \& (\bigoplus_j M_j^2) \\ &\dashv\vdash \bigoplus_i (M_i^1 \& (\bigoplus_j M_j^2)) \\ &\dashv\vdash \bigoplus_{(i,j) \in I \times J} (M_i^1 \& M_j^2) \text{ .} \end{aligned}$$

The case where A has the form $A_1 \otimes A_2$ is exactly analogous, making use instead of the standard $\otimes\text{-}\oplus\text{-distributivity}$ of linear logic. □

\oplus -free fragment above. Then,

$$\Gamma \not\vdash \mathbf{F}$$

and

if $\Gamma \vdash C \oplus D$ then $\Gamma \vdash C$ or $\Gamma \vdash D$.

Proof By cut-elimination any proof of a sequent can be replaced by a cut-free proof. The above lemma follows by induction on the size of cut-free proofs. \square

Lemma 31 For any \multimap -free assertion A ,

$$\llbracket A \rrbracket_N = \{M \mid M \text{ is } \oplus\text{-free, } M \vdash A\} .$$

Proof The proof proceeds by structural induction on the assertion A .

$A \equiv \mathbf{T}$: Clearly, $\llbracket \mathbf{T} \rrbracket_N = \mathcal{M} = \{M \mid M \oplus\text{-free}\} = \{M \oplus\text{-free} \mid M \vdash \mathbf{T}\}$.

$A \equiv \mathbf{F}$: Now, using lemma 30, $\llbracket \mathbf{F} \rrbracket_N = \emptyset = \{M \mid M \oplus\text{-free, } M \vdash \mathbf{F}\}$.

$A \equiv \mathbf{1}$: $\llbracket \mathbf{1} \rrbracket_N = \{M \mid M \rightarrow \mathbf{0}\} = \{M \mid M \oplus\text{-free, } M \vdash \mathbf{1}\}$ by lemma 28.

$A \equiv a$: $\llbracket a \rrbracket_N = \{M \mid M \rightarrow a\} = \{M \mid M \oplus\text{-free, } M \vdash a\}$ by lemma 28.

$A \equiv A_1 \& A_2$: We argue straightforwardly that

$$\begin{aligned} M \in \llbracket A_1 \& A_2 \rrbracket_N &\Leftrightarrow M \in \llbracket A_1 \rrbracket_N \text{ and } M \in \llbracket A_2 \rrbracket_N \text{ by definition,} \\ &\Leftrightarrow M \oplus\text{-free, } M \vdash A_1 \text{ and } M \vdash A_2 \text{ by induction,} \\ &\Leftrightarrow M \oplus\text{-free, } M \vdash A_1 \& A_2 \text{ by the proof rules.} \end{aligned}$$

$$\begin{aligned} A \equiv A_1 \oplus A_2: \text{ Argue: } & M \in \llbracket A_1 \oplus A_2 \rrbracket_N \\ &\Leftrightarrow M \in \llbracket A_1 \rrbracket_N \text{ or } M \in \llbracket A_2 \rrbracket_N \text{ by definition,} \\ &\Leftrightarrow M \oplus\text{-free and either } M \vdash A_1 \text{ or } M \vdash A_2 \text{ by induction,} \\ &\Leftrightarrow M \oplus\text{-free, } M \vdash A_1 \oplus A_2 \text{ by lemma 30 and } (\vdash \oplus). \end{aligned}$$

$A \equiv A_1 \otimes A_2$: The proof in this case is a little more involved. Argue:

$$\begin{aligned} M \in \llbracket A_1 \otimes A_2 \rrbracket_N &\Leftrightarrow \exists M_1 \in \llbracket A_1 \rrbracket_N, M_2 \in \llbracket A_2 \rrbracket_N. M \rightarrow M_1 + M_2 \text{ by definition,} \\ &\Leftrightarrow M \oplus\text{-free, } \exists M_1, M_2 \oplus\text{-free. } M_1 \vdash A_1, M_2 \vdash A_2 \text{ and} \\ &\quad M \vdash M_1 \otimes M_2 \text{ by induction and lemma 28,} \\ &\Rightarrow M \oplus\text{-free, } M \vdash A_1 \otimes A_2 \text{ from the proof rules.} \end{aligned}$$

To show the converse implication, and so equivalence, assume M is \oplus -free and $M \vdash A_1 \otimes A_2$. By lemma 29, we may assume

$$A_1 \dashv\vdash \bigoplus_{i \in I} M_i^1 \text{ and } A_2 \dashv\vdash \bigoplus_{j \in J} M_j^2 .$$

We may furthermore assume I and J to be nonempty. Otherwise $A_1 \otimes A_2 \dashv\vdash \mathbf{F}$ and so, as M is \oplus -free, $M \not\vdash A_1 \otimes A_2$ by lemma 30—a contradiction.

$$M \vdash \left(\bigoplus_{i \in I} M_i^1 \right) \otimes \left(\bigoplus_{j \in J} M_j^2 \right),$$

so by distributivity,

$$M \vdash \bigoplus_{(i,j) \in I \times J} M_i^1 \otimes M_j^2 .$$

Hence, by lemma 30,

$$M \vdash M_i^1 \otimes M_j^2 \text{ for some } i \in I, j \in J$$

such that $M_i^1 \vdash A_1$ and $M_j^2 \vdash A_2$. This plainly gives the required converse. \square

Corollary 32 $\models_N A \text{ iff } \vdash A$, for any \multimap -free assertion A .

Thus we have completeness.

Because we only use the decomposition lemma (lemma 29) for the \otimes case of the structural induction in lemma 31, we also get completeness for the larger fragment of assertions B given by:

$$B ::= A \mid A \multimap B \mid \top \mid \mathbf{F} \mid \mathbf{1} \mid a \mid B_1 \& B_2 \mid B_1 \oplus B_2$$

where A lie in the \multimap -free fragment and a , as usual, ranges over atoms.

Lemma 33 For the larger fragment,

$$\llbracket B \rrbracket_N = \{ M \mid M \oplus\text{-free}, M \vdash B \} .$$

Proof The proof proceeds by structural induction, as in lemma 31, but for a new case where the assertion has the form $A \multimap B$. Because of its assumed form, by lemma 29, there is a decomposition

$$A \dashv\vdash \bigoplus_{i \in I} M_i .$$

Now, for \oplus -free M we argue that

$$\begin{aligned} M \in \llbracket A \multimap B \rrbracket_N &\Leftrightarrow \forall M_A \in \llbracket A \rrbracket_N. M \otimes M_A \in \llbracket B \rrbracket_N \text{ by definition,} \\ &\Leftrightarrow \forall M_A \oplus\text{-free. } M_A \vdash A \Rightarrow M \otimes M_A \vdash B \text{ by induction,} \\ &\Leftrightarrow \forall M_A \oplus\text{-free. } M_A \vdash \bigoplus_{i \in I} M_i \Rightarrow M \otimes M_A \vdash B \\ &\Leftrightarrow \forall M_A \oplus\text{-free. } (\exists i \in I. M_A \vdash M_i) \Rightarrow M \otimes M_A \vdash B \text{ by lemma 30,} \\ &\Leftrightarrow \forall i \in I, \forall M_A \oplus\text{-free. } M_A \vdash M_i \Rightarrow M \otimes M_A \vdash B \\ &\Leftrightarrow \forall i \in I. M \otimes M_i \vdash B. \end{aligned}$$

Here “ \Rightarrow ” follows directly by taking $M_A = M_i$. The converse “ \Leftarrow ” makes use of the fact that if $M_A \vdash M_i$ and $M \otimes M_i \vdash B$ then $M \otimes M_A \vdash B$.

Now, continuing the argument,

$$\begin{aligned}
&\Leftrightarrow \bigoplus_{i \in I} (M \otimes M_i) \vdash B \text{ from the proof system,} \\
&\Leftrightarrow M \otimes (\bigoplus_{i \in I} M_i) \vdash B \\
&\Leftrightarrow M \otimes A \vdash B \\
&\Leftrightarrow M \vdash A \multimap B.
\end{aligned}$$

□

Corollary 34 *For the larger fragment, $\models B$ iff $\vdash B$ with the additional $\&$ - \oplus -distributivity law.*

Theorem 35 *For the \multimap -free fragment,*

$$\Gamma \models A \quad \text{iff} \quad \Gamma \vdash A$$

with the additional $\&$ - \oplus -distributive law.

Proof Corollary 34 gives

$$\models \bigotimes \Gamma \multimap A \text{ iff } \vdash \bigotimes \Gamma \multimap A .$$

Hence

$$\Gamma \models A \text{ iff } \Gamma \vdash A .$$

□

14 Quantification and Atomic Nets

Definition 36 *A net is atomic iff whenever $M \rightarrow \underline{Q}$ then $\underline{Q} \rightarrow M$, for any marking M .*

□

This corresponds to $\mathbf{1}$ being atomic in the associated quantale. A sufficient condition for a net to be atomic is that every transition of it leads to a nonempty multiset of markings.

Notice that $A \& \mathbf{1}$ plays the role of the exponential $!A$, and indeed according the net semantics, when the net N is atomic

$$\llbracket !A \rrbracket_N = \llbracket A \& \mathbf{1} \rrbracket_N .$$

An other interesting consequence of dealing with an atomic net N is, that we no longer need the logical constant, \perp , denoting linear-absurdity in order to express negative properties. Whatever property we could state before in terms of validity of a closed formula A , can now be stated negatively as $\models_N A \& \mathbf{1} \multimap \mathbf{F}$. Precisely:

Proposition 37 *For an atomic net N and a closed formula A ,*

$$\models_N A \& \mathbf{1} \multimap \mathbf{F} \quad \text{iff} \quad \not\models_N A .$$

the denotation of A contains $\mathbf{1}$ and empty otherwise. Hence $\llbracket A \& \mathbf{1} \multimap \mathbf{F} \rrbracket_N$ equals \mathbf{T} in case $\not\models_N A$ and equals \mathbf{F} in case $\models_N A$. \square

Abbreviating $A \& \mathbf{1} \multimap \mathbf{F}$ by $\sim A$, this means that all properties stated in section 10 as A^\perp , now can be stated as $\sim A$. In exercise 49 in section 15 we shall see that \perp in fact can be expressed directly in the extended logic.

Syntax

Assume a countable set of atoms. Define the assertions over the atoms to be:

$$A ::= \mathbf{T} \mid \mathbf{F} \mid \mathbf{1} \mid a \mid x \mid A_1 \otimes A_2 \mid A_1 \multimap A_2 \mid A_1 \& A_2 \mid A_1 \oplus A_2 \mid \bigoplus_x A \mid \big\&_x A$$

where a ranges over the atoms and x ranges over countably many variables. The new constructions $\bigoplus_x A$ and $\big\&_x A$ are forms of existential and universal quantification and bind accordingly. We adopt the traditional notions of free and bound variable and in particular use $\text{FV}(A)$ for the set of free variables in A , and more generally $\text{FV}(A_1, \dots, A_n)$ for $\text{FV}(A_1) \cup \dots \cup \text{FV}(A_n)$. The variables x are to be thought of as standing for markings of a net.

Semantics

Given a net N , with markings (i.e. finite multisets of places) \mathcal{M} , a (marking) environment is a function ρ from variables to markings \mathcal{M} . Because of the presence of free variables we define the meaning of an assertion with respect to a marking environment. In particular,

$$\begin{aligned} \llbracket \bigoplus_x A \rrbracket_N \rho &= \bigcup_{M \in \mathcal{M}} \llbracket A \rrbracket_N \rho[M/x], \\ \llbracket \big\&_x A \rrbracket_N \rho &= \bigcap_{M \in \mathcal{M}} \llbracket A \rrbracket_N \rho[M/x], \\ \llbracket x \rrbracket_N \rho &= \{M \in \mathcal{M} \mid M \rightarrow \rho(x)\}. \end{aligned}$$

Atoms are interpreted as singletons of places of the net as in section 7 and similarly validity of a closed assertion A for the given net, N , can be expressed by:

$$\models_N A \quad \text{iff} \quad \underline{0} \in \llbracket A \rrbracket_N \rho .$$

This is generalized to open terms by taking the universal closure:

$$\models_N A \quad \text{iff} \quad \underline{0} \in \llbracket \big\&_{x_1} \cdots \big\&_{x_k} A \rrbracket_N \rho$$

where A has free variables x_1, \dots, x_k (here ρ can be arbitrary because $\big\&_{x_1} \cdots \big\&_{x_k} A$ is closed).

Let T be a subset of closed assertions in the syntax. Define

$$\models_N (B_1 \otimes \cdots \otimes B_n \multimap A).$$

Before proceeding with the proof rules we show how the new constructions can be used to express liveness.

A transition t is *live* iff $\forall M \in [M_0] \exists M' \in [M]. M' [t]$.

This can be expressed by:

$$\models_N \bigotimes_x ((M_0 \multimap x) \& \mathbf{1} \multimap (x \multimap \bullet t \otimes \top)) .$$

Obviously *liveness* can then be expressed for finite nets.

Exercise 38 In this exercise we consider a logic for Petri nets inspired by branching-time logic. The syntax of the branching formulae is

$$\varphi ::= p \mid \varphi \wedge \varphi \mid \varphi \vee \varphi \mid \neg \varphi \mid \forall G \varphi \mid \exists F \varphi \quad (12)$$

where p range over some atomic propositions. For a net N and a fixed valuation V_N yielding for each atomic proposition a set of markings for which it holds, we can recursively define when a marking M of N satisfies a formula φ , written $M \models_N \varphi$, as follows:

$$\begin{aligned} M \models_N p & \quad \text{iff } M \in V_N(p) \\ M \models_N \varphi_1 \wedge \varphi_2 & \text{iff } M \models_N \varphi_1 \text{ and } M \models_N \varphi_2 \\ M \models_N \varphi_1 \vee \varphi_2 & \text{iff } M \models_N \varphi_1 \text{ or } M \models_N \varphi_2 \\ M \models_N \neg \varphi & \quad \text{iff } M \not\models_N \varphi \\ M \models_N \forall G \varphi & \quad \text{iff } \forall M' \in [M]. M' \models_N \varphi \\ M \models_N \exists F \varphi & \quad \text{iff } \exists M' \in [M]. M' \models_N \varphi . \end{aligned}$$

As an example, a valuation V_N and an atomic proposition p_t could express the set of markings enabling t . I.e. $V_N(p_t) = \{M \mid \exists M'. M [t] M'\}$.

Now, let M be some marking term and X an ordered infinite set of variables such that $\text{FV}(M) \subseteq X$. Furthermore, suppose a set Φ of formulae of (12) for which we, given a marking M and a $\varphi \in \Phi$, have a translation $A_\varphi(M)$ into linear logic, i.e. such that $M \models_N \varphi$ iff $\models_N A_\varphi(M)$. For p_t mentioned above, the translation $A_{\exists F p_t}(M)$ of $\exists F p_t$ could be $M \multimap \bullet t \otimes \top$.

The translation can then be extended to cover all branching formulae over Φ by the following schema

$$\begin{aligned}
\llbracket \varphi \rrbracket_X^M &= \llbracket \varphi \rrbracket_X^M, \text{ for } \varphi \in \Phi \\
\llbracket \varphi_1 \wedge \varphi_2 \rrbracket_X^M &= \llbracket \varphi_1 \rrbracket_X^M \& \llbracket \varphi_2 \rrbracket_X^M \\
\llbracket \varphi_1 \vee \varphi_2 \rrbracket_X^M &= \llbracket \varphi_1 \rrbracket_X^M \oplus \llbracket \varphi_2 \rrbracket_X^M \\
\llbracket \neg \varphi \rrbracket_X^M &= \sim \llbracket \varphi \rrbracket_X^M \\
\llbracket \forall G \varphi \rrbracket_X^M &= \&_x((M \multimap x) \& \mathbf{1} \multimap \llbracket \varphi \rrbracket_{X \cup \{x\}}^x) \\
\llbracket \exists F \varphi \rrbracket_X^M &= \oplus_x((M \multimap x) \& \llbracket \varphi \rrbracket_{X \cup \{x\}}^x)
\end{aligned}$$

where x is the first variable not in X .

This gives another and easy way to see how to express in linear logic that a transition t is live or M -dead because

$$\begin{aligned}
t \text{ live iff } M_0 &\models \forall G \exists F p_t \\
t \text{ is } M\text{-dead iff } M &\models \forall G \neg p_t \text{ iff } M \models \neg \exists F p_t .
\end{aligned}$$

Show that for a *closed* marking term M and branching formula φ over Φ we have:

$$M \models_N \varphi \text{ iff } \models_N \llbracket \varphi \rrbracket_\emptyset^M . \quad \square$$

Proof rules

The proof rules are those of section 2 (without exponentials—they will become definable in the purely propositional logic), together with:

$$\frac{\Gamma \vdash A}{\Gamma[\theta] \vdash A[\theta]} \quad (\text{Subst.})$$

where θ is a substitution of *marking terms* (i.e. assertions built up from variables, atoms and $\mathbf{1}$ purely by \otimes)—the usual care to avoid capture of free variables applies here.

$$\frac{\Gamma, B \vdash A}{\Gamma, \oplus_x B \vdash A} \quad x \notin \text{FV}(\Gamma, A) \quad (\oplus\vdash)$$

$$\frac{\Gamma \vdash A[M/x]}{\Gamma \vdash \oplus_x A} \quad \text{where } M \text{ is a marking term} \quad (\vdash\oplus)$$

Note these rules yield (and in the presence of (Subst.) are equivalent with)

$$\frac{\Gamma, \oplus_x B \vdash A}{\Gamma, B \vdash A} \quad x \notin \text{FV}(\Gamma, A) \quad (\oplus\text{-adj.})$$

Assume $(\oplus\vdash)$ and $(\vdash\oplus)$. The upwards direction of the rule $(\oplus\text{-adj.})$ is simply $(\oplus\vdash)$.

The downwards direction viz.

$$\frac{\Gamma, \oplus_x B \vdash A}{\Gamma, B \vdash A} \quad x \notin \text{FV}(\Gamma, A)$$

$B \vdash \oplus_x B$. Now by the cut rule from the assumption $\Gamma, \oplus_x B \vdash A$ we can conclude $\Gamma, B \vdash A$.

By using (Subst.) we can also derive $(\oplus\vdash)$ and $(\vdash\oplus)$ from $(\oplus\text{-adj.})$. The rule $(\oplus\vdash)$ is simply the upwards reading of $(\oplus\text{-adj.})$. Now we show $(\vdash\oplus)$ follows from $(\oplus\text{-adj.})$: Clearly $\oplus_x A \vdash \oplus_x A$, from which $A \vdash \oplus_x A$ follows by $(\oplus\text{-adj.})$; hence by (Subst.) $A[M/x] \vdash \oplus_x A$, making $(\vdash\oplus)$ derivable.

$$\frac{\Gamma, B \vdash A}{\Gamma, \&_x B \vdash A} \quad (\&\vdash)$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash \&_x A} \quad x \notin \text{FV}(\Gamma) \quad (\vdash\&)$$

Note these rules are equivalent with

$$\frac{\Gamma \vdash A}{\Gamma \vdash \&_x A} \quad x \notin \text{FV}(\Gamma) \quad . \quad (\&\text{-adj.})$$

In addition we have the following axioms valid of nets:

$$(A_1 \oplus A_2) \& B \vdash (A_1 \& B) \oplus (A_2 \& B) \quad (\&\text{-}\oplus\text{-dist.})$$

$$\left(\bigoplus_x A\right) \& B \vdash \bigoplus_x (A \& B) \quad \text{where } x \notin \text{FV}(B) \quad (\&\text{-}\oplus\text{-dist.})$$

In fact, in the presence of the atomicity, basis and primeness axioms, these distributivity laws are derivable from those in the special case where B is $\mathbf{1}$. The other distributive law,

$$\&_x (A \oplus B) \vdash (\&_x A) \oplus B \quad \text{where } x \notin \text{FV}(B),$$

is also derivable (for general B).

$$\vdash (\&_x B) \oplus \bigoplus_x ((B \& \mathbf{1}) \multimap \mathbf{F}) \quad (\text{Atomicity})$$

These entail sequents of the following form (by taking the variable x to not appear in B):

$$\vdash B \oplus ((B \& \mathbf{1}) \multimap \mathbf{F}) \quad .$$

These hold because in an atomic net the denotation of a formula $B \& \mathbf{1}$, in an environment for its free variables, only has two possibilities, to be the denotation of \mathbf{F} or the denotation of $\mathbf{1}$.

$$A \vdash \bigoplus_x x \otimes ((x \multimap A) \& \mathbf{1}) \quad \text{where } x \notin \text{FV}(A) \quad (\text{Basis})$$

These hold in an atomic net because an assertion denotes a set of markings; notice how the expression $(x \multimap A) \& \mathbf{1}$ is equivalent to $\mathbf{1}$ in the case the marking

a contribution of x when it satisfies A .

$$\begin{aligned}
& (x \multimap \mathbf{F}) \vdash \mathbf{F} \\
& (x \multimap B \oplus C) \vdash (x \multimap B) \oplus (x \multimap C) \quad (\text{Primeness}) \\
& (x \multimap \oplus_y A) \vdash \oplus_y (x \multimap A) \text{ where } y \text{ and } x \text{ are distinct.}
\end{aligned}$$

These axioms hold because if a marking is contained in union, denoting a disjunction, then it is clearly in a component of the union.

For clarity we have collected the new proof rules:

$\frac{\Gamma \vdash A}{\Gamma[\theta] \vdash A[\theta]} \quad (\text{Subst.})$	
$\frac{\Gamma \vdash A[M/x]}{\Gamma \vdash \oplus_x A} \quad (\vdash\oplus)$	$\frac{\Gamma, B \vdash A}{\Gamma, \oplus_x B \vdash A} \quad x \notin \text{FV}(\Gamma, A) \quad (\oplus\vdash)$
$\frac{\Gamma \vdash A}{\Gamma \vdash \&_x A} \quad x \notin \text{FV}(\Gamma) \quad (\vdash\&)$	$\frac{\Gamma, B \vdash A}{\Gamma, \&_x B \vdash A} \quad (\&\vdash)$
$(A_1 \oplus A_2) \& B \vdash (A_1 \& B) \oplus (A_2 \& B) \quad (\&\oplus\text{-dist.})$	
$(\oplus_x A) \& B \vdash \oplus_x (A \& B), \quad x \notin \text{FV}(B) \quad (\&\oplus\text{-dist.})$	
$\vdash (\&_x B) \oplus \oplus_x ((B \& \mathbf{1}) \multimap \mathbf{F}) \quad (\text{Atomicity})$	
$A \vdash \oplus_x x \otimes ((x \multimap A) \& \mathbf{1}), \quad x \notin \text{FV}(A) \quad (\text{Basis})$	
$(x \multimap \mathbf{F}) \vdash \mathbf{F}$	
$(x \multimap B \oplus C) \vdash (x \multimap B) \oplus (x \multimap C) \quad (\text{Primeness})$	
$(x \multimap \oplus_y A) \vdash \oplus_y (x \multimap A) \text{ where } y \text{ and } x \text{ are distinct}$	

The soundness of the basis and atomicity axioms follows from the fact that, in an atomic net,

$$\llbracket A \& \mathbf{1} \rrbracket_{N\rho} = \begin{cases} \mathbf{1} & \text{if } \mathbf{1} \subseteq \llbracket A \rrbracket_{N\rho} \\ \mathbf{F} & \text{otherwise} \end{cases} \quad \text{and} \quad \llbracket A \& \mathbf{1} \multimap \mathbf{F} \rrbracket_{N\rho} = \begin{cases} \mathbf{F} & \text{if } \mathbf{1} \subseteq \llbracket A \rrbracket_{N\rho} \\ \mathbf{T} & \text{otherwise.} \end{cases}$$

We have already remarked that in an atomic net, an exponential $!A$ is represented by $A \& \mathbf{1}$. In fact from the atomicity axioms and the rules for exponentials, there is a fairly direct proof of their equivalence, yielding

$$!A \dashv\vdash A \& \mathbf{1} \quad (13)$$

—the syntax of exponentials can be eliminated in favour of the purely propositional connectives.

$\&$. From $A \& \mathbf{1} \vdash (A \& \mathbf{1}) \& \mathbf{1}$ and (Atomicity) we also get $\vdash (A \& \mathbf{1}) \oplus (A \& \mathbf{1} \multimap \mathbf{F})$. Using $A \& \mathbf{1} \vdash A \& \mathbf{1}$ we then deduce

$$A \& \mathbf{1} \vdash ((A \& \mathbf{1}) \otimes (A \& \mathbf{1})) \oplus ((A \& \mathbf{1}) \otimes (A \& \mathbf{1} \multimap \mathbf{F}))$$

so that $A \& \mathbf{1} \vdash ((A \& \mathbf{1}) \otimes (A \& \mathbf{1})) \oplus \mathbf{F}$, i.e.

$$A \& \mathbf{1} \vdash (A \& \mathbf{1}) \otimes (A \& \mathbf{1}) .$$

By (4) of section 3 this ensures $A \& \mathbf{1} \vdash !A$, and clearly $!A \vdash A \& \mathbf{1}$, making (13).

Completeness

Prime theories play a key role in showing completeness. A prime theory arises as the set of all assertions true of some net, and conversely from a prime theory we can construct a net whose truths are precisely the prime theory. The completeness proof proceeds by showing that if an assertion is not provable then there is a prime theory, and so a net, for which it fails to hold.

In constructing prime theories we follow Henkin and extend the original syntax to include new atoms drawn from

$$c_0, c_1, \dots, c_n, \dots$$

a countably infinite enumeration of atoms not already present in the syntax. Suppose C is a subset of $\{c_i \mid i \in \omega\}$. Suppose Γ, A are assertions from the syntax extended by C , and that \mathcal{F} is a theory (i.e. a subset of assertions) of the extended syntax. We use

$$\Gamma \vdash_{\mathcal{F}}^C A$$

to mean the sequent is provable in the proof system for the extended syntax, using the assertions in \mathcal{F} as axioms. A judgment $\Gamma \vdash A$ means a sequent is provable in the proof system of the original assertion language, without extra atoms.

Lemma 39 *Let B be a closed assertion and \mathcal{F} a theory in a syntax extended by atoms $C \subseteq \{c_i \mid i \in \omega\}$. Then*

$$\Gamma \vdash_{\mathcal{F} \cup \{B\}}^C A \quad \text{iff} \quad \Gamma, B \& \mathbf{1} \vdash_{\mathcal{F}}^C A .$$

Proof “ \Rightarrow ” By induction on the size of derivations of $\Gamma \vdash_{\mathcal{F} \cup \{B\}}^C A$, considering the final rule used.

“ \Leftarrow ” From $\vdash_{\mathcal{F} \cup \{B\}} B$ and $\vdash \mathbf{1}$ follows $\vdash_{\mathcal{F} \cup \{B\}} B \& \mathbf{1}$ and so the results from hypothesis and cut. \square

syntax extended by atoms C , is called a prime theory iff

- (i) $F \notin \mathcal{F}$,
 $A_1 \oplus A_2 \in \mathcal{F} \Rightarrow A_1 \in \mathcal{F}$ or $A_2 \in \mathcal{F}$,
 $\oplus_x A \in \mathcal{F} \Rightarrow A[M/x] \in \mathcal{F}$, for some (necessarily closed) marking term M .
- (ii) \mathcal{F} is deductively closed, i.e. A closed and $\vdash_{\mathcal{F}}^C A \Rightarrow A \in \mathcal{F}$. □

Lemma 41 (Existence of prime theories). *Let A be an assertion and T a subset of closed assertions in the original syntax, for which*

$$\not\vdash_T A .$$

Then, there is a prime theory \mathcal{F} , consisting of assertions over the syntax extended by some $C \subseteq \{c_i \mid i \in \omega\}$, such that

$$T \subseteq \mathcal{F} \text{ and } \not\vdash_{\mathcal{F}}^C A .$$

Proof As the atoms and variables form countable sets we can enumerate all the assertions

$$A_0, A_1, \dots, A_n, \dots$$

of the syntax extended by atoms $\{c_i \mid i \in \omega\}$.

By induction, for $n \in \omega$, we define a chain of deductively – closed theories \mathcal{F}_n with new atoms C_n , such that

$$T \subseteq \mathcal{F}_n \text{ and } \not\vdash_{\mathcal{F}_n}^{C_n} A .$$

Take $\mathcal{F}_0 = \{B \mid \vdash_T B\}$. Clearly $T \subseteq \mathcal{F}_0$ and, by assumption, $\not\vdash_{\mathcal{F}_0} A$.

Assuming \mathcal{F}_n is deductively closed, includes T , and $\not\vdash_{\mathcal{F}_n}^{C_n} A$, define \mathcal{F}_{n+1} according to the following cases:

- (i) n is even, and there is an assertion $B_1 \oplus B_2 \in \mathcal{F}_n$ with $B_1 \notin \mathcal{F}_n, B_2 \notin \mathcal{F}_n$.
- (ii) n is odd, and there is an assertion $\oplus_x B \in \mathcal{F}_n$ with $B[M/x] \notin \mathcal{F}_n$ for any closed marking terms M in the syntax over atoms extended by C_n .
- (iii) neither (i) nor (ii) applies.

In case (iii), define $C_{n+1} = C_n$ and $\mathcal{F}_{n+1} = \mathcal{F}_n$.

In case (i), take the earliest assertion in the enumeration $B_1 \oplus B_2 \in \mathcal{F}_n$ and $B_1 \notin \mathcal{F}_n$ and $B_2 \notin \mathcal{F}_n$. As \mathcal{F}_n is deductively-closed, $(B_1 \oplus B_2) \& \mathbf{1} \in \mathcal{F}_n$, so

$$(B_1 \& \mathbf{1}) \oplus (B_2 \& \mathbf{1}) \in \mathcal{F}_n$$

$$\vdash_{\mathcal{F}_n, B_1}^{C_n} A \text{ and } \vdash_{\mathcal{F}_n, B_2}^{C_n} A .$$

Then, by lemma 39,

$$B_1 \& \mathbf{1} \vdash_{\mathcal{F}_n}^{C_n} A \text{ and } B_2 \& \mathbf{1} \vdash_{\mathcal{F}_n}^{C_n} A .$$

Hence $(B_1 \& \mathbf{1}) \oplus (B_2 \& \mathbf{1}) \vdash_{\mathcal{F}_n}^{C_n} A$. But this implies $\vdash_{\mathcal{F}_n}^{C_n} A$, a contradiction. Thus

$$\not\vdash_{\mathcal{F}_n, B_1}^{C_n} A \text{ or } \not\vdash_{\mathcal{F}_n, B_2}^{C_n} A .$$

Supposing, for instance, $\not\vdash_{\mathcal{F}_n, B_1}^{C_n} A$, take \mathcal{F}_{n+1} to be $\{D \text{ closed} \mid \vdash_{\mathcal{F}_n, B_1}^{C_n} D\}$ and $C_{n+1} = C_n$.

In case (ii), take the earliest, according to the enumeration, $\bigoplus_x B \in \mathcal{F}_n$ for which $B[M/x] \notin \mathcal{F}_n$ for all marking terms M , and where x is not a free variable of A . As \mathcal{F}_n is deductively-closed, $(\bigoplus_x B) \& \mathbf{1} \in \mathcal{F}_n$, so

$$\bigoplus_x (B \& \mathbf{1}) \in \mathcal{F}_n .$$

Let c be the first new atom in the list c_0, c_1, \dots which is not in C_n . Define $C_{n+1} = C_n \cup \{c\}$ and \mathcal{F}_{n+1} to consist of all closed assertions in the deductive closure of $\mathcal{F}_n \cup \{B[c/x]\}$, i.e.

$$\mathcal{F}_{n+1} = \{D \text{ closed} \mid \vdash_{\mathcal{F}_n, B[c/x]}^{C_{n+1}} D\} .$$

We must check that $\not\vdash_{\mathcal{F}_{n+1}}^{C_{n+1}} A$. To this end, assume otherwise, that $\vdash_{\mathcal{F}_n, B[c/x]}^{C_{n+1}} A$. Then, by lemma 39,

$$B[c/x] \& \mathbf{1} \vdash_{\mathcal{F}_n}^{C_{n+1}} A .$$

As c does not appear in C_n or \mathcal{F}_n ,

$$B \& \mathbf{1} \vdash_{\mathcal{F}_n}^{C_n} A .$$

To obtain the proof of this sequent, replace all occurrences of the new atom c in the proof of $B[c/x] \& \mathbf{1} \vdash_{\mathcal{F}_n}^{C_{n+1}} A$ by a new variable—one which does not appear anywhere in the proof—and finally use (Subst.) to replace this variable by x using the fact that renaming bound variables preserves logical equivalence. But now we can deduce

$$\bigoplus_x (B \& \mathbf{1}) \vdash_{\mathcal{F}_n}^{C_n} A .$$

But $\bigoplus_x (B \& \mathbf{1}) \in \mathcal{F}_n$ and \mathcal{F}_n is deductively-closed making $\vdash_{\mathcal{F}_n}^{C_n} A$, a contradiction. Thus $\not\vdash_{\mathcal{F}_{n+1}}^{C_{n+1}} A$, as required.

In this way, we inductively define a chain of theories \mathcal{F}_n over the syntax extended by C_n , such that

$$C_n \subseteq C_{n+1} \text{ and } \mathcal{F}_n \subseteq \mathcal{F}_{n+1},$$

with $T \subseteq \mathcal{F}_0$. Finally take $C = \bigcup_{n \in \omega} C_n$ and $\mathcal{F} = \bigcup_{n \in \omega} \mathcal{F}_n$ to form the required prime theory. \square

by taking:

- *Places* to be the original atoms, including those of C . Atoms are interpreted as their singleton markings.
- *Transitions* as those pairs (M, M') of multisets of places for which $\widehat{M} \vdash_{\mathcal{F}}^C \widehat{M}'$.

We use \mathcal{M} to represent the set of all markings of the net N . (Note the markings coincide with the closed marking terms of \mathcal{F} .)

Lemma 42 *For markings M, M' in the net N ,*

$$M \rightarrow M' \text{ in } N \quad \text{iff} \quad M \vdash_{\mathcal{F}}^C M' .$$

Furthermore N is atomic.

Proof The proof of “iff” is like that for lemma 23.

Regarding atomicity of the net, suppose $M \rightarrow \underline{0}$, for $M \in \mathcal{M}$. Then $M \vdash_{\mathcal{F}}^C \mathbf{1}$. Certainly

$$M \oplus (M \& \mathbf{1} \multimap \mathbf{F}) \in \mathcal{F}$$

and as \mathcal{F} is a prime theory

$$M \in \mathcal{F} \text{ or } M \& \mathbf{1} \multimap \mathbf{F} \in \mathcal{F} .$$

The case $M \& \mathbf{1} \multimap \mathbf{F} \in \mathcal{F}$ is impossible. To see this assume otherwise, that $M \& \mathbf{1} \multimap \mathbf{F} \in \mathcal{F}$. As $M \vdash_{\mathcal{F}}^C \mathbf{1}$, and clearly $M \vdash M$, it follows that $M \vdash_{\mathcal{F}}^C M \& \mathbf{1}$ and hence $M \vdash_{\mathcal{F}}^C \mathbf{F}$. This ensures $\vdash_{\mathcal{F}}^C (M \multimap \mathbf{F})$ which together with the primeness axiom

$$(M \multimap \mathbf{F}) \vdash \mathbf{F}$$

yields $\vdash_{\mathcal{F}}^C \mathbf{F}$. I.e. $\mathbf{F} \in \mathcal{F}$ —contradicting \mathcal{F} being a prime theory. The case $M \& \mathbf{1} \multimap \mathbf{F} \in \mathcal{F}$ is thus impossible.

$M \in \mathcal{F}$, where $M \vdash_{\mathcal{F}}^C \mathbf{1}$. Thus $M \rightarrow \underline{0}$ in the net. The net N is therefore atomic. \square

We need the following facts:

Lemma 43

- (i) *Let B be an assertion with $\text{FV}(B) \subseteq \{x\}$. Let the assertions Γ not include x as a free variable. Then*

$$(\forall M \in \mathcal{M}. \Gamma \vdash_{\mathcal{F}}^C B[M/x]) \Rightarrow \Gamma \vdash_{\mathcal{F}}^C \&_x B .$$

which x_1, \dots, x_k are not free. Then

$$(\forall M_1, \dots, M_k \in \mathcal{M}. \Gamma \vdash_{\mathcal{F}}^C B[M_1/x_1, \dots, M_k/x_k]) \Rightarrow \Gamma \vdash_{\mathcal{F}}^C B .$$

Proof (i) We first prove the special case

$$(\forall M \in \mathcal{M}. \vdash_{\mathcal{F}}^C A[M/x]) \Rightarrow \vdash_{\mathcal{F}}^C \&_x A,$$

where $\text{FV}(A) \subseteq \{x\}$, by contraposition. By atomicity,

$$\vdash (\&_x A) \oplus \bigoplus_x (A \& \mathbf{1} \multimap \mathbf{F}),$$

where both operands of \oplus are closed, so as \mathcal{F} is a prime theory

$$\vdash_{\mathcal{F}}^C \&_x A \text{ or } \vdash_{\mathcal{F}}^C \bigoplus_x (A \& \mathbf{1} \multimap \mathbf{F}) .$$

Thus supposing $\not\vdash_{\mathcal{F}}^C \&_x A$ we obtain $\vdash_{\mathcal{F}}^C \bigoplus_x (A \& \mathbf{1} \multimap \mathbf{F})$. But as \mathcal{F} is a prime theory, there is then $M \in \mathcal{M}$, such that $\vdash_{\mathcal{F}}^C (A[M/x] \& \mathbf{1} \multimap \mathbf{F})$ and thus $\not\vdash_{\mathcal{F}}^C A[M/x]$.

The more general statement of (i) above follows by taking $A \equiv \otimes \Gamma \multimap B$ where Γ does not contain x free and B has at most x free. Note $\Gamma \vdash \&_x B$ follows from $\vdash \&_x (\otimes \Gamma \multimap B)$.

(ii) follows by induction using (i) together with the fact that $\&_{x_k} B \vdash B$. \square

Lemma 44 For assertions Γ , B and A suppose $\Gamma, B[M/x] \vdash_{\mathcal{F}}^C A$ for all $M \in \mathcal{M}$. Assume $\text{FV}(B) \subseteq \{x\}$. Then

$$\Gamma, \bigoplus_x B \vdash_{\mathcal{F}}^C A .$$

Proof We show that if $B[M/x] \vdash_{\mathcal{F}}^C A$ for all $M \in \mathcal{M}$ where $\text{FV}(B) \subseteq x$, then $\bigoplus_x B \vdash_{\mathcal{F}}^C A$. The seemingly stronger result follows by \multimap -adjunction.

Assume $B[M/x] \vdash_{\mathcal{F}}^C A$, for all $M \in \mathcal{M}$, and $\text{FV}(B) \subseteq \{x\}$. Let some $M \in \mathcal{M}$ be given. Since $\text{FV}(B[M/x]) \neq \emptyset$ (M is closed) we can use $(\vdash \&)$ to get $B[M/x] \vdash_{\mathcal{F}}^C A'$ where $A' \equiv \&_{y \in \text{FV}(A)} A$. I.e. $\text{FV}(A') = \emptyset$ and $\vdash_{\mathcal{F}}^C B[M/x] \multimap A'$ for all $M \in \mathcal{M}$. By (i) of lemma 43 we then get

$$\vdash_{\mathcal{F}}^C \&_x (B \multimap A')$$

and by logic that

$$\vdash_{\mathcal{F}}^C (\bigoplus_x B) \multimap A' .$$

Here we used the logic deduction:

$$\begin{aligned}
&\Rightarrow \&_x(B \multimap A'), B \vdash A' \\
&\Rightarrow \&_x(B \multimap A'), \oplus_x B \vdash A' \quad \text{by } (\oplus\vdash) \text{ as } x \notin \text{FV}(\&_x(B \multimap A'), A'), \\
&\Rightarrow \&_x(B \multimap A') \vdash (\oplus_x B) \multimap A'.
\end{aligned}$$

From $\vdash_{\mathcal{F}}^C (\oplus_x B) \multimap A'$ we get $\oplus_x B \vdash_{\mathcal{F}}^C A'$ and so $\oplus_x B \vdash_{\mathcal{F}}^C A$ by ($\&$ -adj.). \square

Now we can relate semantics in the net N to provability in the prime theory \mathcal{F} :

Lemma 45 *For any assertion A , for any marking environment ρ ,*

$$\llbracket A \rrbracket_N \rho = \{M \in \mathcal{M} \mid M \vdash_{\mathcal{F}}^C A[\rho]\} .$$

Proof By structural induction on A .

$$A \equiv \top: \llbracket \top \rrbracket_N \rho = \mathcal{M} = \{M \in \mathcal{M} \mid M \vdash_{\mathcal{F}}^C \top\}.$$

$$A \equiv \text{F}: \llbracket \text{F} \rrbracket_N \rho = \emptyset = \{M \in \mathcal{M} \mid M \vdash_{\mathcal{F}}^C \text{F}\} \text{---as } \text{F} \notin \mathcal{F} \text{ and } (M \multimap \text{F}) \vdash \text{F}.$$

$$A \equiv \mathbf{1}: \llbracket \mathbf{1} \rrbracket_N \rho = \{M \in \mathcal{M} \mid M \rightarrow \mathbf{0}\} = \{M \in \mathcal{M} \mid M \vdash_{\mathcal{F}}^C \mathbf{1}\} \text{ by lemma 42.}$$

$$A \equiv a: \llbracket a \rrbracket_N \rho = \{M \in \mathcal{M} \mid M \rightarrow a\} = \{M \in \mathcal{M} \mid M \vdash_{\mathcal{F}}^C a\} \text{ by lemma 42.}$$

$$A \equiv x: \llbracket x \rrbracket_N \rho = \{M \in \mathcal{M} \mid M \rightarrow \rho(x)\} = \{M \in \mathcal{M} \mid M \vdash_{\mathcal{F}}^C \rho(x)\} \text{ by lemma 42.}$$

$A \equiv A_1 \otimes A_2$: For $M \in \mathcal{M}$,

$$\begin{aligned}
&M \in \llbracket A_1 \otimes A_2 \rrbracket_N \rho \\
&\Leftrightarrow \exists M_1 \in \llbracket A_1 \rrbracket_N \rho, M_2 \in \llbracket A_2 \rrbracket_N \rho. M \rightarrow M_1 + M_2, \\
&\Leftrightarrow \exists M_1, M_2 \in \mathcal{M}. M_1 \vdash_{\mathcal{F}}^C A_1[\rho], M_2 \vdash_{\mathcal{F}}^C A_2[\rho] \text{ and } M \vdash_{\mathcal{F}}^C M_1 \otimes M_2, \\
&\Rightarrow M \vdash_{\mathcal{F}}^C (A_1 \otimes A_2)[\rho].
\end{aligned}$$

To show “ \Leftarrow ” as well, we write $A'_1 \equiv A_1[\rho]$ and $A'_2 \equiv A_2[\rho]$, and use the basic facts that

$$A'_1 \vdash \oplus_{x_1} x_1 \otimes ((x_1 \multimap A'_1) \& \mathbf{1}) \quad \text{and} \quad A'_2 \vdash \oplus_{x_2} x_2 \otimes ((x_2 \multimap A'_2) \& \mathbf{1}) .$$

Assuming $M \vdash_{\mathcal{F}}^C A'_1 \otimes A'_2$, we obtain

$$M \vdash_{\mathcal{F}}^C \left(\bigoplus_{x_1} x_1 \otimes ((x_1 \multimap A'_1) \& \mathbf{1}) \right) \otimes \left(\bigoplus_{x_2} x_2 \otimes ((x_2 \multimap A'_2) \& \mathbf{1}) \right) .$$

By distributivity of \otimes over \oplus ,

$$M \vdash_{\mathcal{F}}^C \bigoplus_{x_1} \bigoplus_{x_2} x_1 \otimes ((x_1 \multimap A'_1) \& \mathbf{1}) \otimes x_2 \otimes ((x_2 \multimap A'_2) \& \mathbf{1}) .$$

By primeness and because \mathcal{F} is a prime theory,

$$M \vdash_{\mathcal{F}}^C M_1 \otimes ((M_1 \multimap A'_1) \& \mathbf{1}) \otimes M_2 \otimes ((M_2 \multimap A'_2) \& \mathbf{1}) \quad (*)$$

By atomicity and the fact that \mathcal{F} is a prime theory

$$\begin{aligned} & \vdash_{\mathcal{F}}^C (M_1 \multimap A'_1) \quad \text{or} \quad \vdash_{\mathcal{F}}^C ((M_1 \multimap A'_1) \& \mathbf{1} \multimap \mathbf{F}) \quad \text{and} \\ & \vdash_{\mathcal{F}}^C (M_2 \multimap A'_2) \quad \text{or} \quad \vdash_{\mathcal{F}}^C ((M_2 \multimap A'_2) \& \mathbf{1} \multimap \mathbf{F}) . \end{aligned}$$

As $M \not\vdash_{\mathcal{F}}^C \mathbf{F}$ and $M_i \not\vdash_{\mathcal{F}}^C \mathbf{F}$ for $i = 1, 2$ (by the same argument as in the $A \equiv \mathbf{F}$ case above) we from (*) deduce $(M_i \multimap A'_i) \& \mathbf{1} \not\vdash_{\mathcal{F}}^C \mathbf{F}$ for $i = 1, 2$, and we must have $\vdash_{\mathcal{F}}^C (M_1 \multimap A'_1)$ and $\vdash_{\mathcal{F}}^C (M_2 \multimap A'_2)$, i.e. $M_1 \vdash_{\mathcal{F}}^C A'_1$, and $M_2 \vdash_{\mathcal{F}}^C A'_2$, as well as $M \vdash_{\mathcal{F}}^C M_1 \otimes M_2$. This establishes the required reverse implication, and so equivalence.

$$\begin{aligned} A \equiv A_1 \multimap A_2: \quad & M \in \llbracket A_1 \multimap A_2 \rrbracket_N \rho \\ & \Leftrightarrow \forall M_1 \in \llbracket A_1 \rrbracket_N \rho. M + M_1 \in \llbracket A_2 \rrbracket_N \rho. \\ & \Leftrightarrow \forall M_1 \vdash_{\mathcal{F}} A_1[\rho]. M \otimes M_1 \vdash_{\mathcal{F}}^C A_2[\rho] \text{ by induction,} \\ & \Leftrightarrow \forall M_1 \in \mathcal{M}. M \otimes M_1 \otimes ((M_1 \multimap A_1[\rho]) \& \mathbf{1}) \vdash_{\mathcal{F}}^C A_2[\rho] \end{aligned}$$

where the last equivalence relies on atomicity and the fact that \mathcal{F} is a prime theory. In more detail, writing $A'_1 \equiv A_1[\rho]$, $A'_2 \equiv A_2[\rho]$, we have

$$(i) \quad \vdash_{\mathcal{F}}^C M_1 \multimap A'_1 \quad \text{or} \quad (ii) \quad \vdash_{\mathcal{F}}^C (M_1 \multimap A'_1) \& \mathbf{1} \multimap \mathbf{F}$$

for any $M_1 \in \mathcal{M}$. For case (i), $(M_1 \multimap A'_1) \& \mathbf{1} \not\vdash_{\mathcal{F}}^C \mathbf{1}$. In case (ii), $(M_1 \multimap A'_1) \& \mathbf{1} \not\vdash_{\mathcal{F}}^C \mathbf{F}$.

It follows, by considering the two cases, that for any $M \in \mathcal{M}'$

$$(M_1 \vdash_{\mathcal{F}}^C A'_1 \Rightarrow M \otimes M_1 \vdash_{\mathcal{F}}^C A'_2) \text{ iff } M \otimes M_1 \otimes ((M_1 \multimap A'_1) \& \mathbf{1}) \vdash_{\mathcal{F}}^C A'_2 .$$

Now note that by lemma 44,

$$\begin{aligned} & \forall M_1 \in \mathcal{M}. M \otimes M_1 \otimes ((M_1 \multimap A'_1) \& \mathbf{1}) \vdash_{\mathcal{F}}^C A'_2 \\ & \Leftrightarrow M \otimes \oplus_{x_1} x_1 \otimes ((x_1 \multimap A'_1) \& \mathbf{1}) \vdash_{\mathcal{F}}^C A'_2, \\ & \Leftrightarrow M \otimes A'_1 \vdash_{\mathcal{F}}^C A'_2 \text{ as } A'_1 \not\vdash_{\mathcal{F}}^C \oplus_{x_1} x_1 \otimes ((x_1 \multimap A'_1) \& \mathbf{1}), \\ & \Leftrightarrow M \vdash_{\mathcal{F}}^C A'_1 \multimap A'_2 \text{ i.e. } M \vdash_{\mathcal{F}}^C A_1[\rho] \multimap A_2[\rho], \text{ as required.} \end{aligned}$$

$$\begin{aligned} A \equiv A_1 \oplus A_2: \quad & M \in \llbracket A_1 \oplus A_2 \rrbracket_N \rho \Leftrightarrow M \in \llbracket A_1 \rrbracket_N \rho \text{ or } M \in \llbracket A_2 \rrbracket_N \rho \\ & \Leftrightarrow M \vdash_{\mathcal{F}}^C A_1[\rho] \text{ or } M \vdash_{\mathcal{F}}^C A_2[\rho] \\ & \Leftrightarrow M \vdash_{\mathcal{F}}^C A_1[\rho] \oplus A_2[\rho]. \end{aligned}$$

where primeness and the fact that \mathcal{F} is a prime theory is used in showing “ \Leftarrow ”; the other direction “ \Rightarrow ” follows directly from the proof system.

$$\begin{aligned} A \equiv A_1 \& A_2: \quad & M \in \llbracket A_1 \& A_2 \rrbracket_N \rho \Leftrightarrow M \in \llbracket A_1 \rrbracket_N \rho \text{ and } M \in \llbracket A_2 \rrbracket_N \rho \\ & \Leftrightarrow M \vdash_{\mathcal{F}}^C A_1[\rho] \text{ and } M \vdash_{\mathcal{F}}^C A_2[\rho] \text{ by induction,} \\ & \Leftrightarrow M \vdash_{\mathcal{F}}^C A_1[\rho] \& A_2[\rho]. \end{aligned}$$

$$\begin{aligned} A \equiv \oplus_x B : \quad & M \in \llbracket \oplus_x B \rrbracket_N \rho \Leftrightarrow \exists M' \in \mathcal{M}. M \in \llbracket B \rrbracket_N \rho[M'/x] \\ & \Leftrightarrow \exists M' \in \mathcal{M}. M \vdash_{\mathcal{F}}^C B[\rho[M'/x]] \text{ by induction,} \\ & \Leftrightarrow M \vdash_{\mathcal{F}}^C (\oplus_x^4 B)[\rho]. \end{aligned}$$

ness and \mathcal{F} being a prime theory.

$$\begin{aligned}
A \equiv \&x B: M \in \llbracket \&x B \rrbracket_N \rho &\Leftrightarrow \forall M' \in \mathcal{M}. M \in \llbracket B \rrbracket_N \rho[M'/x] \\
&&\Leftrightarrow \forall M' \in \mathcal{M}. M \vdash_{\mathcal{F}}^C B[\rho[M'/x]] \text{ by induction,} \\
&&\Leftrightarrow M \vdash_{\mathcal{F}}^C (\&x B)[\rho] \text{ by (i) of lemma 43.}
\end{aligned}$$

□

Theorem 46 (Completeness).

Let A, B and $\Gamma \equiv B_1, \dots, B_n$ be assertions and T consist of closed assertions in the original syntax. Then,

$$\Gamma \models_T A \quad \text{iff} \quad \Gamma \vdash_T A .$$

Proof As \multimap is present as a constructor on assertions it is clearly sufficient to show

$$\models_T A \text{ iff } \vdash_T A .$$

The “if” direction is shown by induction on the proof of $\vdash_T A$. To show the “only if” direction we prove its contraposition:

$$\not\vdash_T A \Rightarrow \not\models_T A .$$

Suppose A is an assertion with free variables x_1, \dots, x_k . Suppose $\not\vdash_T A$. Then there is a prime theory $\mathcal{F} \supseteq T$ over additional constants C such that $\not\vdash_{\mathcal{F}}^C A$. Let N be the net constructed from \mathcal{F} ; let \mathcal{M} be the set consisting of its markings. As we now show, $\not\models_N A$. Suppose otherwise, that $\models_N A$. Recall

$$\llbracket A \rrbracket_N \rho = \{M \in \mathcal{M} \mid M \vdash_{\mathcal{F}}^C A[\rho]\}$$

by lemma 45. Hence as $\models_N A$ means $\underline{\mathcal{Q}} \in \llbracket A \rrbracket_N \rho$ for all environments ρ , we see

$$\vdash_{\mathcal{F}}^C A[\rho] \text{ for all environments } \rho .$$

Therefore $\vdash_{\mathcal{F}}^C A[M_1/x_1, \dots, M_k/x_k]$ for all $M_1, \dots, M_k \in \mathcal{M}$. Hence by (ii) of lemma 43, $\vdash_{\mathcal{F}}^C A$, a contradiction. Thus N is an atomic net satisfying all axioms of the theory T and yet $\not\models_N A$. Hence $\not\models_T A$, as required. □

Exercise 47 (Infinite disjunctions)

An alternative complete proof system with respect to countable atomic nets can be obtained by adding a countably infinite disjunction and infinitary rules for reasoning about it. More precisely, as the syntax of assertions take

$$A ::= \mathbf{T} \mid \mathbf{F} \mid \mathbf{1} \mid a \mid A_1 \otimes A_2 \mid A_1 \multimap A_2 \mid A_1 \& A_2 \mid \bigoplus_{i \in I} A_i,$$

where a ranges over a countable set of atoms and I over countable indexing sets drawn from ω (this avoids our syntax becoming a proper class). When $I = \{1, 2\}$,

semantics of assertions in an atomic net N is like that earlier in section 7, with the additional assumption that every place is named by an atomic proposition (this assumption ensures that we can describe all markings within the logic of assertions). The semantics of countable disjunctions $\bigoplus_{i \in I} A_i$ is a slight extension of that in the finite case:

$$\llbracket \bigoplus_{i \in I} A_i \rrbracket_N = \bigcup_{i \in I} \llbracket A_i \rrbracket_N$$

All other semantic clauses are the same as in section 7. As proof rules we take those of earlier, in section 2, but replacing the rules for binary \oplus by the following two proof rules, the second of which is infinitary:

$$\frac{\Gamma \vdash A_j}{\Gamma \vdash \bigoplus_{i \in I} A_i} \text{ for } j \in I, \quad \frac{\Gamma, B_i \vdash C \text{ for all } i \in I}{\Gamma, \bigoplus_{i \in I} B_i \vdash C}.$$

In addition to these logical rules we also have rules specific to atomic nets:

$$\begin{aligned} & (\bigoplus_{i \in I} A_i) \& B \vdash \bigoplus_{i \in I} (A_i \& B) \\ & \vdash B \oplus (B \& \mathbf{1} \multimap \mathbf{F}) \\ & A \vdash \bigoplus_{M \in \mathcal{M}} M \otimes ((M \multimap A) \& \mathbf{1}) \\ & M \multimap \bigoplus_{i \in I} A_i \vdash \bigoplus_{i \in I} (M \multimap A_i) \end{aligned}$$

where $M \in \mathcal{M}$. Here \mathcal{M} denotes the set of finite multisets of atoms. As earlier, in the context of a assertions, we make the identification with such multisets and tensor formulae. Note that \mathcal{M} is countable as the set of atoms is assumed countable. Note, for instance, that the last rule amounts to

$$(M \multimap \mathbf{F}) \vdash \mathbf{F}, \text{ for } M \in \mathcal{M},$$

when the indexing set I is empty.

Show that the proof system described above is sound and complete with respect to atomic nets.

[Hint: Mimic the completeness proof above, taking as a prime theory a deductively closed subset of assertions with the property that if it contains a disjunction then it contains a disjunct. To show that any theory not entailing A can be extended to a prime theory, define a suitable chain of theories \mathcal{F}_n by transfinite induction on ordinals n , assuming inductively that \mathcal{F}_n is deductively-closed and $A \notin \mathcal{F}_n$, and take the required prime theory as their union.] \square

15 Decidability Issues

Here we show that the question of whether an assertion from the logic of section 14 belongs to the theory of a net is undecidable. We use the following proposition.

$$\langle x \mid A \rangle \equiv \bigoplus_x [(A \ \& \ 1) \otimes x]$$

for an assertion A and variable x . Its denotation with respect to a net valuation N and environment ρ is given by:

$$\llbracket \langle x \mid A \rangle \rrbracket_{N\rho} = \downarrow \{M \mid \underline{Q} \in \llbracket A \rrbracket_{N\rho}[M/x]\} .$$

Exercise 49

- (i) Prove proposition 48 above.
- (ii) Show that linear absurdity \perp (cf. section 10) is definable for atomic nets, in the following way:

$$\perp \equiv \langle x \mid x \ \& \ 1 \ \multimap \mathbf{F} \rangle$$

i.e., for an arbitrary environment,


$$\llbracket \langle x \mid x \ \& \ 1 \ \multimap \mathbf{F} \rangle \rrbracket_{N\rho} = \{M \mid \underline{Q} \not\vdash M \text{ in } N\} .$$

- (iii) Show that for an atomic net N ,

$$\llbracket \langle x \mid \sim(M_0 \multimap x) \rangle \rrbracket_{N\rho} = \{M \mid M_0 \not\vdash M \text{ in } N\} . \quad \square$$

Theorem 50 *The property $\models_N A$ is undecidable for a finite net N and assertion A of the logic of section 14 (assumed to have infinitely many atomic propositions).*

Proof From Rabin's work (described in [Hac76]), we know that the question of whether the reachable markings of one finite net are included in the reachable markings of another is undecidable. More precisely, for two finite nets N_1 and N_2 , over the same set of places, with markings M_1, M_2 , it is undecidable whether or not all the markings reachable from M_1 in N_1 are included in those reachable from M_2 in N_2 . If $\models_N A$ were decidable we could decide Rabin's problem. From N_1, N_2 we effectively construct another net N as follows: Its places are those in common to N_1 and N_2 plus two additional, distinct places a_1 and a_2 . Its transitions are obtained as a disjoint union of those of N_1 and N_2 ; we adjoin a_1 (with multiplicity 1) to the pre and post places of transitions originally in N_1 and a_2 to the pre and post places of those transitions originally of N_2 :



Now, using proposition 48, we could decide Rabin's problem if we could decide

$$\models_N \langle x \mid \sim(M_2 \otimes a_2 \multimap x \otimes a_2) \rangle \multimap \langle x \mid \sim(M_1 \otimes a_1 \multimap x \otimes a_1) \rangle$$

Hence the property $\models_N A$ is undecidable. \square

Exercise 51 For this exercise suppose we modify $\models_N A$ to be as before so that not only does every atomic proposition name a place of N , but also every place in N is named by some atomic proposition of the logic of section 14 (so every place of N receives a name).

- (i) Show that with this modification we can *not* achieve strong completeness:

$$\models_T A \text{ iff } \vdash_T A$$

for an arbitrary theory T . Proceed as follows: For a finite net N the set of assertions

$$T_N = \{M \multimap M' \mid M \rightarrow M' \text{ in } N\} \cup \{\sim(M \multimap M') \mid M \not\rightarrow M' \text{ in } N\}$$

is recursively enumerable (see [May84]). Show that with our modified understanding of satisfaction

$$\models_N A \text{ iff } \models_{T_N} A.$$

Deduce that there is no effectively given sound and *strongly* complete proof system.

- (ii) On the other hand, assuming the atomic propositions form an infinite set, show that there is a complete proof system in the sense that

$$\vdash A \text{ iff } \models A$$

where the proof rules are those of section 14.

[Hint: Its proof involves a slight modification of the notion of a prime theory.]

\square

Although we feel sure the logic of section 14 is undecidable i.e. $\vdash A$ is not decidable for general A , we presently lack a proof (unfortunately, the techniques of [LMSN90] do not work directly because of our primeness axioms). There remain, in addition, the questions of decidability and complexity of weaker fragments of the logic than that of section 14 and their theories with respect to a specified finite net. However, directly from [LMSN90], it is known that the \oplus -free and \multimap -free fragments (without the law of $\&$ - \oplus -distributivity) of section 13 are decidable.

Two questions emerge in our work:

- (i) What can Petri nets contribute to linear logic?
- (ii) What can linear logic contribute to Petri nets?

The expressiveness and completeness results that follow give answers to (i). For the \oplus -free fragment of linear logic (its syntax is given in Section 13), Petri nets are a good model; they give a sound and complete interpretation of this fragment of the logic without having to add any extra axioms or rules. For the other parts of linear logic that we treat, the answer must be more qualified, because we need to adjoin extra axioms to obtain completeness. Addressing (i), we hope the reader comes to share our view that Petri nets (with our semantics) are a helpful, intuitive and convincing model of linear logic—certainly Petri nets carry their own intuition and are models of interest, independently of linear logic.

Answering (ii) is harder at this stage. Linear logic is one of few logics for Petri nets which treat in a non-artificial way the fact that places can hold with multiplicities. The logic of section 14 is also very expressive and we have a complete proof system, although we pay through the theory of a net being non-recursive (see theorem 50). On the negative side, it's sometimes very hard to see if and how a particular property can be expressed, though this criticism could also be levelled at the very useful modal μ -calculus and so cannot be taken too seriously. A fuller answer to (ii) would rest on a more detailed investigation, including that of fragments and variants of linear logic and whether they are useful in the study of Petri nets.⁴

In summary, it is claimed that this paper helps in the understanding of linear logic. The paper exposes a possible role linear logic may have in reasoning about Petri nets and, through them, in concurrent computation.

⁴One possible irritation with the present semantics is that all atoms must name places, or more precisely their downwards closure with respect to reachability, possibly in a many-one way. Gian Luca Cattani has shown in [Cat] that the semantics can be relaxed to allow atoms to denote false at the slight cost of adding existence conditions to the logic and changing the substitution rule. Another extension would be to add transition names to the logic.

Answer to exercise 1 The proofs are:

$$(i) \quad \frac{\Gamma \vdash A \& B \quad \frac{\overline{A \vdash A}}{A \& B \vdash A}}{\Gamma \vdash A} \quad \text{and} \quad \frac{\frac{\overline{A \vdash A} \quad \overline{B \vdash B}}{A, B \vdash A \otimes B} \quad \Gamma, A \otimes B \vdash C}{\Gamma, A, B \vdash C}$$

$$(ii) \quad \frac{\frac{\frac{\overline{A \vdash A}}{A \vdash A \oplus B} \quad \overline{C \vdash C}}{A, (A \oplus B) \multimap C \vdash C} \quad \frac{\frac{\overline{B \vdash B}}{B \vdash A \oplus B} \quad \overline{C \vdash C}}{B, (A \oplus B) \multimap C \vdash C}}{\frac{(A \oplus B) \multimap C \vdash A \multimap C \quad (A \oplus B) \multimap C \vdash B \multimap C}{(A \oplus B) \multimap C \vdash (A \multimap C) \& (B \multimap C)}}$$

$$(iii) \quad \frac{\frac{\frac{\overline{A \vdash A} \quad \overline{B \vdash B}}{A, B \vdash A \otimes B} \quad \frac{\overline{A \vdash A} \quad \overline{C \vdash C}}{A, C \vdash A \otimes C}}{A, B \vdash (A \otimes B) \oplus (A \otimes C) \quad A, C \vdash (A \otimes B) \oplus (A \otimes C)}}{\frac{A, (B \oplus C) \vdash (A \otimes B) \oplus (A \otimes C)}{A \otimes (B \oplus C) \vdash (A \otimes B) \oplus (A \otimes C)}} \quad \text{and}$$

$$\frac{\frac{\frac{\overline{A \vdash A} \quad \overline{B \vdash B}}{A, B \vdash A \otimes (B \oplus C)} \quad \frac{\overline{C \vdash C}}{C \vdash B \oplus C}}{A, C \vdash A \otimes (B \oplus C)} \quad \frac{\overline{A \vdash A} \quad \overline{C \vdash C}}{A, C \vdash A \otimes (B \oplus C)}}{\frac{A \otimes B \vdash A \otimes (B \oplus C) \quad A \otimes C \vdash A \otimes (B \oplus C)}{(A \otimes B) \oplus (A \otimes C) \vdash A \otimes (B \oplus C)}}$$

Answer to exercise 4

$$(i) \quad p \leq q \Leftrightarrow p \vee q = q \Leftrightarrow r \otimes (p \vee q) = r \otimes q \Leftrightarrow r \otimes p \vee r \otimes q = r \otimes q \Leftrightarrow r \otimes p \leq r \otimes q.$$

$$(ii) \quad \text{only if: } r \otimes p \leq q \Rightarrow r \leq \vee\{r' \mid r' \otimes p \leq q\} = p \multimap q.$$

if: Notice that

$$(p \multimap q) \otimes p = (\vee\{r \mid r \otimes p \leq q\}) \otimes p = \vee\{r \otimes p \mid r \otimes p \leq q\} \leq q$$

and hence by monotonicity $r \leq p \multimap q \Rightarrow r \otimes p \leq (p \multimap q) \otimes p \leq q$.

Answer to exercise 7

$$\llbracket d \rrbracket_N = \{\underline{d}, \underline{a} + \underline{b}, \underline{c}\}$$

$$\llbracket d \otimes e \rrbracket_N = \{\underline{d} + \underline{e}, \underline{a} + \underline{b} + \underline{c}, 2\underline{c}, \underline{a} + \underline{b} + \underline{e}, \underline{d} + \underline{c}, \underline{c} + \underline{e}\}$$

$$\llbracket d \& e \rrbracket_N = \{\underline{c}\}$$

- [AD93] G. T. Allwein and J. M. Dunn. Kripke models of linear logic. *The Journal of Symbolic Logic*, 58(2):514–545, 1993.
- [AJ92] Samson Abramsky and R. Jagadeesan. Games and full completeness for multiplicative linear logic. In Rudrapatna Shyamasundar, editor, *FST and TCS 12, Foundations of Software Technology and Theoretical Computer Science, New Delhi, India, December 18–20*, pages 291–301. Springer-Verlag (*LNCS*, 652), 1992.
- [Asp87] Andrea Asperti. A Logic for Concurrency. manuscript, November 1987.
- [AV88] Samson Abramsky and Steve Vickers. Linear Process Logic. Notes by Steve Vickers, 1988.
- [Ber78] G. Berry. Stable models of typed λ -calculi. In *Fifth International Colloquium on Automata, Languages and Programs*, pages 72–89. Springer-Verlag (*LNCS* 62), 1978.
- [Bla92] A. Blass. A game semantics for linear logic. *Annals of Pure and Applied Logic*, 56:183–220, 1992.
- [Bro89] Carolyn Brown. Relating Petri Nets to Formulae of Linear Logic. Technical Report ECS LFCS 89-87, University of Edinburgh, 1989.
- [Cat] Gian Luca Cattani. An existence predicate for a linear logic of Petri net.
- [CGW89] T. Coquand, C. Gunter, and G. Winskel. Domain theoretic models of polymorphism. *Information and Computation*, 81(2), 1989.
- [EW90] Uffe Henrik Engberg and Glynn Winskel. Petri Nets as Models of Linear Logic. In *CAAP '90, Coll. on Trees in Algebra and Programming Copenhagen, Denmark, May 15–18*, pages 147–161. Springer-Verlag (*LNCS* 431), 1990. Appears as Technical Report, DAIMI PB–301.
- [EW93] Uffe Henrik Engberg and Glynn Winskel. Completeness Results for Linear Logic on Petri Nets (Extended Abstract). In *MFCS '93, Mathematical Foundations of Computer Science, Gdańsk, Poland, August 30 - September 3*. Springer-Verlag (*LNCS* 711), 1993. Appears as Technical Report, DAIMI PB–435.
- [GG89a] Carl Gunter and Vijay Gehlot. A Proof-Theoretic Operational Semantics for True Concurrency. Preliminary Report, 1989.

- Report MS-CIS-89-68, University of Pennsylvania, October 1989.
- [Gir86] Jean-Yves Girard. *The system F of variable types, fifteen years later*. *Theoretical Computer Science*, 45, 1986.
- [Gir87] Jean-Yves Girard. Linear Logic. *Theoretical Computer Science*, 50(1):1–102, 1987.
- [GL87] Jean-Yves Girard and Yves Lafont. Linear Logic and Lazy Computation. In *Proc. TAPSOFT 87 (Pisa), vol. 2*, pages 52–66. Springer-Verlag (*LNCS* 250), 1987.
- [Hac76] M. H. T. Hack. *Decidability questions for Petri nets*. PhD thesis, MIT, 1976.
- [Laf88] Yves Lafont. The Linear Abstract Machine. *Theoretical Computer Science*, 59:157–180, 1988.
- [LMSN90] P. Lincoln, J. Mitchell, A. Scedrov, and Shankar N. Decision problems for propositional linear logic. In *Foundations of Computer Science (FOCS'90), volume II*, pages 662–671, St. Louis, MO, October 1990.
- [May84] E. W. Mayr. An algorithm for the general Petri net reachability problem. *SIAM Journal of Computing*, 13(3):441–459, 1984.
- [MOM89] Narciso Martí-Oliet and José Meseguer. From Petri Nets to Linear Logic. In *Category Theory and Computer Science, Manchester, UK*. Springer-Verlag (*LNCS* 389), 1989.
- [MOM91] Narciso Martí-Oliet and José Meseguer. From Petri Nets to Linear Logic: a Survey. *International Journal of Foundations of Computer Science*, 2(4):297–399, 1991.
- [Rei85] Wolfgang Reisig. *Petri Nets, An Introduction*, volume 4 of *EATCS Monographs on Theoretical Computer Science*. Springer-Verlag, 1985.
- [Ros89] Kimmo I. Rosenthal. A Note on Girard Quantales. To appear in: *Cah. de Top. et G. D.*, 1989.
- [Sam] G. Sambin. . Manuscript reported to us by Per Martin-Löf.
- [Yet] D. Yetter. Quantales and Non-Commutative Linear Logic. (preprint).

Recent Publications in the BRICS Report Series

- RS-94-1** Glynn Winskel. *Semantics, Algorithmics and Logic: Basic Research in Computer Science. BRICS Inaugural Talk.* February 1994, 8 pp.
- RS-94-2** Alexander E. Andreev. *Complexity of Nondeterministic Functions.* February 1994, 47 pp.
- RS-94-3** Uffe H. Engberg and Glynn Winskel. *Linear Logic on Petri Nets.* February 1994, 54 pp.
- RS-94-4** Nils Klarlund and Michael I. Schwartzbach. *Graphs and Decidable Transductions based on Edge Constraints.* February 1994, 19 pp. Appears in: *Trees in Algebra and Programming CAAP '94* (ed. S. Tison), LNCS 787, 1994.
- RS-94-5** Peter D. Mosses. *Unified Algebras and Abstract Syntax.* March 1994, 21 pp. To appear in: *Recent Trends in Data Type Specification* (ed. F. Orejas), LNCS 785, 1994.
- RS-94-6** Mogens Nielsen and Christian Clausen. *Bisimulations, Games and Logic.* April 1994, 37 pp. Full version of paper to appear in: *New Results and Trends in Computer Science*, LNCS, 1994.