


PARALLEL CONTEXT-FREE LANGUAGES

by
Sven Skyum

DAIMI PB-30

June 1974

Institute of Mathematics University of Aarhus
DEPARTMENT OF COMPUTER SCIENCE
Ny Munkegade - 8000 Aarhus C - Denmark
Phone 06-12 83 55



PARALLEL CONTEXT-FREE LANGUAGES

Abstract

The relation between the family of context - free languages and the family of parallel context - free languages is examined in this paper.

It is proved that the families are incomparable.

Finally we prove that the family of languages of finite index is contained in the family of parallel context - free languages.

Introduction

It has been an open problem whether or not the family of context – free languages is contained in the family of parallel context – free languages(E. g. ,cf. Rosenfeld (1971)). By "parallel" we mean that whenever you use a production $A \rightarrow \alpha$ in a derivation you have to use this specific production simultaneously for all occurrences of A in the sentential form you are dealing with.

In Siromoney and Krithivasan (to appear) it is claimed that the intersection of the two families consists of languages of finite index.

The reason for writing this paper is that the proofs in Siromoney and Krithivasan (to appear) contain so many serious gaps that it seems appropriate to find alternative proofs.

In this paper it is proved that the family of parallel context – free languages contains the family of languages of finite index but not the family of context – free languages.

It is assumed that the reader is familiar with the basic notions concerning formal language theory. For unexplained notions we refer to Salomaa (1973).

The relation between parallel context-free languages and context-free languages.

Definition

Let $G = (V, T, P, S)$ be an ordinary context-free grammar.

The parallel direct yield relation $\Rightarrow_G^{(p)}$ on the set $(V \cup T)^*$ is defined as follows:

$\alpha \xrightarrow{(p)}_G \beta$ iff

$\alpha = \alpha_1 A \alpha_2 A \dots A \alpha_k$ and $\beta = \alpha_1 w \alpha_2 w \dots w \alpha_k$, where

$\alpha_i \in ((V \cup T) \setminus \{A\})^*$ for $1 \leq i \leq k$ and $A \rightarrow w$ is a production in G . The relation $\Rightarrow_G^{(p)*}$ is the reflexive transitive closure of $\Rightarrow_G^{(p)}$. $\alpha \Rightarrow_G^{(p)t} \beta$ iff

there exists words $\alpha_i \in (V \cup T)^*$ for $0 \leq i \leq t$ such that

$\alpha = \alpha_0 \xrightarrow{(p)}_G \alpha_1 \xrightarrow{(p)}_G \alpha_2 \xrightarrow{(p)}_G \dots \xrightarrow{(p)}_G \alpha_t = \beta$.

The language generated in parallel by G is defined by:

$$L_p(G) = \{ w \in T^* \mid S \Rightarrow_G^{(p)*} w \}$$

The family of parallel context-free languages is the family of languages generated in parallel by ordinary context-free grammars.

The following example shows that the family of parallel context-free languages contains languages which are non-context-free.

Example

The language $\{ a^{2^n} \mid n \geq 0 \}$ is generated in parallel by the following grammar. $(\{S\}, \{a\}, \{S \rightarrow SS, S \rightarrow a\}, S)$.

Theorem 1

The family of context-free languages is not contained in the family of parallel context-free languages.

Proof

Let L be the context-free language generated by the following grammar G .

$G = (\{S, A\}, \{0, 1, (,), [,]\}, P, S)$ where P consists of the following productions.

$S \rightarrow SS, S \rightarrow (0A1], S \rightarrow \lambda,$

$A \rightarrow 0A1, A \rightarrow)S[.$

L is very similar to the Dyck-set (over 0 and 1), but in L you have that corresponding groups of 0's and 1's are surrounded by parantheses and brackets resp.

Assume that there exists a grammar

$H = (V, T, P, S)$ such that H generates L in parallel.

Let t be the number of nonterminals in H and m the length of the longest righthandside of a production in H . Let $\bar{m} = \lceil \log_2 m \rceil + 1$ (the least integer which is greater than or equal to $\log_2 m$ plus one).

We assume that no nonterminal is useless in H and whenever nothing else is stated \Rightarrow and \Rightarrow^* are used instead of $\xRightarrow{(p)}_H$ and $\xRightarrow{(p)^*}_H$.

Lemma 1

If $A \in V$ and $A \Rightarrow^* w_1 w_2$ then $\#_1(w_1) \leq \#_0(w_1) + m^{2t}$ and $\#_0(w_2) \leq \#_1(w_2) + m^{2t}$.

Proof

Since A is not useless there exists a derivation of the form

$S \xRightarrow{t^I} \alpha_1 A \alpha_2 \xRightarrow{t^{II}} v_1 A v_2 A \dots A v_k \xRightarrow{*} v_1 w_1 w_2 v_2 \dots w_1 w_2 v_k \in L,$
 where $t^I, t^{II} \leq t$, $\alpha_i \in (V \cup T)^*$ for $1 \leq i \leq 2$, $v_i \in T^*$ for $1 \leq i \leq k$,
 $w_1 w_2 \in T^*$, and $A \Rightarrow^* w_1 w_2$. Therefore $|v_i| \leq m^{2t}$ for $1 \leq i \leq k$.

As for the Dyck set (over 0 and 1) we have that in any prefix of a word in the language the number of 1's in the prefix is at most the number of 0's.

We have $\#_1(v_1 w_1) \leq \#_0(v_1 w_1)$ which implies that $\#_1(w_1) \leq \#_0(w_1) + |v_1| \leq \#_0 + m^{2^t}$. The same kind of argument used on the suffix $w_2 v_k$ gives that $\#_0(w_2) \leq \#_1(w_2) + m^{2^t}$.

Define inductively Q_i to be the languages.

$Q_1 = \{(0^n)[1^n] \mid n \geq 1\}$, and for $i \geq 1$

$Q_{i+1} = \{(0^n) w_1 w_2 [1^n] \mid n \geq 1; w_1, w_2 \in Q_i\}$.

We have $Q_i \subseteq L$ for all $i \geq 1$.

Let $w \in Q_q$ be the word

$(0^{n_1})(0^{n_2}) \dots (0^{n_q})[1^{n_q}](0^{n_{q+1}}) \dots [1^{n_2}](0^{n_{2^{q-1}+1}})(0^{n_{2^{q-1}+2}}) \dots$
 $\dots [1^{n_{2^{q-1}+1}}][1^{n_1}]$, where $q = (t+2)\bar{m}$, $n_1 = m^{2^t} + 1$,

and $n_i = n_{i-1} + 1$, $2 \leq i \leq 2^q - 1$. w will be fixed in the following and let

(1) $S = w_0 \Rightarrow w_1 \Rightarrow w_2 \Rightarrow \dots \Rightarrow w_n = w$ be a fixed derivation of w in H .

Lemma 2

If A occurs in w_i for some $0 \leq i \leq n$ and this occurrence of A derives the subword w_A of w in the derivation (1), then there exists integers $-2 \leq j \leq 0$ and $0 \leq k$ such that $\#_j(w_A) = 2^k + j$. We will say that A in that case is of degree k , if $k \geq 2$.

Proof

Let w_A be as above and let $h: T^* \rightarrow \{a, b\}^*$ be the homomorphism:

$h(_) = a$, $h(_) = b$, and $h(0) = h(1) = h(_) = h(_) = \lambda$.

Let $v_i \in \{a, b\}^*$ be defined inductively by:

$$v_0 = \lambda$$

$$v_{i+1} = a v_i v_i b, \text{ for } i \geq 0.$$

We then have that $h(w) = v_q$, and $\#_a(v_k) = 2^k - 1$.

If $h(w) = x \underline{a} v_i v_i \underline{b} y$ for some i and $x, y \in \{a, b\}^*$, then we say that the underlined occurrences of a and b correspond to each other, and we will use expressions as "the occurrence of b corresponding to the occurrence of a " in this case.

Now if $|h(w_A)| \leq 1$ then $0 \leq \#_a(h(w_A)) = \#_b(w_A) \leq 1$ which is of the form $2^k + j$ for some $k \geq 0$ and $-2 \leq j \leq 0$.

Assume then that $|h(w_A)| > 1$.

There are then 4 cases to consider, namely:

$$(1) h(w_A) = \dots av_a \quad \text{for some } v \in \{a, b\}^*$$

$$(2) h(w_A) = \dots av_b \quad \text{for some } v \in \{a, b\}^*$$

$$(3) h(w_A) = \dots bva \quad \text{for some } v \in \{a, b\}^*$$

$$(4) h(w_A) = \dots bvb \quad \text{for some } v \in \{a, b\}^*$$

Cases (1), (3), and (4) are all treated in the same way.

Assume e. g. that $h(w_A) = \dots av_a$ for some $v \in \{a, b\}^*$, then $w_A = x(0^p)z$ where $\#_1(z) = 0$ and $p > m^{2^t}$ which is contradictory to Lemma 1.

Assume then that $h(w_A) = \dots av_b$ for some $v \in \{a, b\}^*$. There are again more possibilities:

- (1) The occurrence of b in $h(w)$ which corresponds to the first occurrence of a in the subword $h(w_A)$ is outside $h(w_A)$. The situation is then that the last occurrence of b in $h(w_A)$ must correspond to an occurrence of a inside $h(w_A)$. We then have that $h(w_A) = av^i av_i v_i b$

for some $0 \leq i \leq q$ and $v' \in \{a, b\}^*$.

$$(Ia) \ v' = v''a$$

Then $w_A = x(o^{p_1})(o^{p_2})y_1y_2z$ for some $x, z \in T^*$ and $y_1, y_2 \in Q_i$. $\#_0((o^{p_1})(o^{p_2})y_1y_2z) - \#_1((o^{p_1})(o^{p_2})y_1y_2z) = p_1 + p_2 + \#_0(z) - \#_1(z) \geq p_1 + \#_0(z) > m^{2t}$ which is contradictory to Lemma 1.

$$(Ib) \ v' = v''b$$

Then $h(w_A) = av_iv_ibav_iv_ib$ and $\#_j(w_A) = 2^{i+2} - 2$ or

$$h(w_A) = a\bar{v}av_iv_ibav_iv_ib \text{ for some } 0 \leq i \leq q \text{ and } \bar{v} \in \{a, b\}^*$$

$$(Iba) \ \bar{v} = \bar{v}a$$

The situation is then as in (Ia).

$$(Ibb) \ \bar{v} = \bar{v}b$$

This can not be the case because $bav_iv_ibav_iv_ib$ can not be a subword of $h(w)$.

$$(Ibc) \ \bar{v} = \lambda$$

Then $h(w_A) = aav_iv_ibav_iv_ib$ and $\#_j(w_A) = 2^{i+2} - 1$.

$$(Ic) \ v' = \lambda$$

Then $h(w_A) = aav_iv_ib$ and $\#_j(w_A) = 2^{i+1}$.

(II) The occurrence of b corresponding to the first occurrence of a in $h(w_A)$ is inside $h(w_A)$. We then have that $h(w_A) = av_iv_ibv'b$ for some $0 \leq i \leq q$ and $v' \in \{a, b\}^*$, or $h(w_A) = av_iv_ib$ and $\#_j(w_A) = 2^{i+1} - 1$. The former situation is quite analogous to the situation in (I).

Let $(\alpha_2, \alpha_3, \dots, \alpha_q)$ be a $q-1$ -tuple of functions mapping the words w_j into integers. $\alpha_k(w_i)$ is the number of occurrences of symbols in w_i which are of degree k .

We have then

$\alpha_k(w_0) = 1$ for $k=q$ and 0 otherwise, and

$\alpha_k(w_n) = 0$ for all k .

Assume that $k > \bar{m}$ and $\alpha_k(w_{i+1}) = \alpha_k(w_i) - p$ for some $0 \leq i \leq q-1$ and $p > 0$. Then $\sum_{r=1}^{\bar{m}} \alpha_{k-r}(w_{i+1}) \geq \sum_{r=1}^{\bar{m}} \alpha_{k-r}(w_i) + 2p$.

To see that we observe that in $i+1$ 'th step in the derivation (1) a non-terminal A of degree k must be rewritten as α containing no nonterminal of degree k , but containing at least two occurrences of nonterminals of degree at least $k - \bar{m}$ and at most $k-1$. This is because α has to generate at least $2^k - 2$ rightparantheses and that the length of α is at most m .

Finally we then conclude that there must exist an integer $0 \leq i \leq n$

such that $\sum_{r=0}^{(t+1)\bar{m}} \alpha_{q-r}(w_i) \geq t + 1$.

This means that there in w_i is more than t occurrences of nonterminals generating at least 2 rightparantheses in w . This contradicts the fact that there are only t nonterminals and that the number of 0's between left- and rightparantheses all are different in the word w .

We have then proved that L is not a parallel context-free language.

Theorem 2

The family of context-free language of finite index is contained in the family of parallel context-free languages.

Proof

Let L be a context-free language of index k and $G = (V, T, P, S)$ a context-free grammar of index k generating G .

Define $G' = (V', T, P', S')$ to be the grammar where

$$T' = A, P' = \{ A \rightarrow V, 1 \leq i \leq k \},$$

$$V' = \{ A[i] \mid A \in V, 1 \leq i \leq k \},$$

P' contains all productions of the form

$$A[i_0] \rightarrow x_1 A_1[i_1] x_2 A_2[i_2] \dots A_n[i_n] x_{n+1} \text{ where } x_j \in T^* \text{ for}$$

$1 \leq i_j \leq n+1, 1 \leq i_j \leq k \text{ for all } 0 \leq j \leq n, \text{ and } A_1, A_2, \dots, A_n \in V.$

$A \rightarrow x_1 A_1 x_2 A_2 \dots A_n x_{n+1}$ is a production in P .

$$S' = S[1].$$

It is now obvious that $L(G) = L(G')$, and that G' is of index k . This means that we can choose derivations such that no nonterminal in V' occurs more than once in every sentential form in these derivations. It is then clear that G' generates L also in parallel.

References

Ginsburg, S. and Spanier, E. (1968), Derivation - bounded languages. J. Comput. System Sci. 2, 228-250.

Rosenfeld, A. (1971), Isotonic Grammars, Parallel Grammars, and Picture Grammars. Machine Intelligence 6, 281-294.

Salomaa, A. (1973), Formal Languages. Academic Press.

Siromoney, R. and Krithivasan, K. (to appear), Parallel context-free languages. Information and Control.